

Certificate of Mailing

Date of Deposit September 18, 2001

Label Number: EL714747653US

I hereby certify under 37 C.F.R. § 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" with sufficient postage on the date indicated above and is addressed to BOX PATENT APPLICATION, Assistant Commissioner for Patents, Washington, D.C. 20231.

Guy Beardsley

Printed name of person mailing correspondence


Signature of person mailing correspondence

APPLICATION
FOR
UNITED STATES LETTERS PATENT

APPLICANTS : Pierre Michetti, Irene Corthesy-Theulaz, Andre Blum,
Catherine Davin, Rainer Haas, Jean-Pierre Kraehenbuhl,
Emilia Saraga

TITLE : Urease-Based Vaccine and Treatment for Helicobacter
Infection

0955739-091801
FOR 60-6E/55660


S P E C I F I C A T I O N

UREASE-BASED VACCINE AND TREATMENT FOR HELICOBACTER INFECTION

The present invention relates to the prevention and treatment of gastric infection in mammals, including humans.

5 More particularly, the present invention relates to a vaccine suitable for use in the prevention and treatment of Helicobacter infection in mammals, including humans, and to a method of treatment of humans suffering from gastric infection, its consequences such as chronic gastritis or peptic ulcer, and prevention of gastric cancer.4

RELATED APPLICATIONS

 This application is a continuation-in-part of U.S. application Serial No. 08/085,938, filed July 6, 1993, which is a continuation-in-part application of U.S. application Serial No. 07/970,996, filed November 3, 1992, the whole of which applications (including drawings) are hereby incorporated by reference.

085739-091801
10

15

BACKGROUND OF THE INVENTION

Helicobacter infections of human gastric epithelium cause gastritis, are a major factor in the development of peptic ulcers and gastric lymphoma, and may be a risk factor for the development of gastric cancer. (Blaser, M.J. "Gastric Campylobacter-like Organisms, Gastritis and Peptic Ulcer Disease" Gastroenterology, vol. 93, 371-383 (1987); Graham, D.Y. "Campylobacter pylori and Peptic Ulcer Disease" Gastroenterology, vol. 196, 615-625 (1989); Parsonnet, J. et al. "Helicobacter pylori Infection in Intestinal and Diffuse-Type Gastric Adenocarcinomas" J. Natl. Cancer Inst., vol. 93, 640-643 (1991); Wotherspoon, A.C., et al., "Regression of Primary Low-Grade B-Cell Gastric Lymphoma of Mucosa-Associated Lymphoid Tissue Type After Eradication of Helicobacter pylori," Lancet, vol. 342, 575-577 (1993)). The most frequent infection agent is Helicobacter pylori, followed at a much lower frequency by Helicobacter heilmanii. Helicobacter pylori is a slender S-shaped gram negative microorganism, which is routinely recovered from gastric biopsies of adults and children with histologic evidence of gastritis or peptic ulceration. Evidence for a causal relationship between Helicobacter pylori and gastroduodenal disease comes from studies in human volunteers, patients with ulcers and gastric cancer, gnotobiotic pigs, and germ-free rodents. Regarding etiology, Koch's postulates were satisfied by creating

histologically confirmed gastritis in previously uninfected individuals following consumption of viable microorganisms.

(Marshall, B.J. et al. "Attempt to Fulfill Koch's Postulate for pyloric Campylobacter" Med. J. Aust., vol. 142, 436-439 (1985);

5 Morris, A. et al. "Ingestion of Campylobacter pyloritis Causes Gastritis and Raised Fasting Gastric pH" Am. J. Gastroenterol., vol. 82, 192-199 (1987); Engstrand, L. et al. "Inoculation of Barrier-Born Pigs With Helicobacter pylori: A Useful Animal Model for Gastritis Type B" Infect. Immun., vol. 53, 1763-1768 (1990); Fox, J.G. et al. "Gastric Colonization by Campylobacter pylori Subsp. mustelae in Ferrets" Infect. Immun., vol. 56, 2994-2996 (1988); Fox, J.G. et al. "Helicobacter mustelae-Associated Gastritis in Ferrets: An Animal Model of Helicobacter pylori Gastritis in Humans" Gastroenterology, vol. 99, 352-361 (1990);
10 Lee, A. et al. "A Small Animal Model of Human Helicobacter pylori Active Chronic Gastritis" Gastroenterology, vol. 99, 1315-1323 (1990); Fox, J.G. et al. "Helicobacter Felis Gastritis in Gnotobiotic Rats: An Animal Model of Helicobacter pylori Gastritis" Infect. Immun., vol. 59, 785-791 (1991); Eaton, K.A.
15 et al. "Campylobacter pylori Virulence Factors in Gnotobiotic Piglets" Infect. Immun., vol. 57, 1119-1125 (1989)), and by treatment to eradicate Helicobacter pylori, with resolution of the gastritis and, in patients with peptic ulcer disease, a decrease in the recurrence rate. (Peterson, W.L. "Helicobacter

pylori and Peptic Ulcer Disease" N. Engl. J. Med., vol. 324,
1043-1048 (1991)).

5 Gastroduodenal diseases thought to be associated with
Helicobacter infection include acute, chronic, and atrophic
gastritis, peptic ulcer disease including both gastric and
duodenal ulcers, gastric cancer, chronic dyspepsia with severe
erosive gastroduodenitis, refractory non-ulcer dyspepsia,
intestinal metaplasia, and low grade MALT lymphoma. Helicobacter
infection is also the principle cause of asymptomatic chronic
gastritis.

10 In spite of in vitro susceptibility to many
antimicrobial agents, in vivo eradication of established
Helicobacter pylori infections with antimicrobial agents is often
difficult to achieve. (Czinn, S.J. and Nedrud, J.G. "Oral
Immunization Against Helicobacter pylori" Infect. Immun., vol.
59, 2359-2363 (1991)). The microorganism is found within the
mucous coat overlying the gastric epithelium and in gastric pits.
These are locations which do not appear to allow for adequate
antimicrobial levels to be achieved even when antibiotics are
20 given orally at high doses. At the present time, most
authorities recommend a "triple therapy", namely a bismuth salt
in combination with drugs such as tetracycline and metronidazole
for 2-4 weeks. However, the effectiveness of this or other
chemotherapeutic regimens remains suboptimal. Recently, a

109055739-091801
108T60" 5E25560

National Institutes of Health panel of medical experts recommended a triple therapy with bismuth, tetracycline and metronidazole, administered for two weeks for treatment of peptic ulcers (Cimons, M., "Drug Combination Found Effective on Peptic
5 Ulcers," L.A. Times at A14 (Feb. 10, 1994)). However, this treatment is commonly associated with diarrhea and it may produce serious adverse drug reactions. (See, Dick-Hegedus, E. and Lee, A., "Use of a Mouse Model to Examine Anti-Helicobacter pylori Agents," Scand. J. Gastroenterol., vol. 26, 909-915 (1991)).
10 Treatment with antibiotics also may not solve the problem of reinfection and there is evidence for a high incidence of reinfection in some studies (Coelho, L.G., et al., "Duodenal Ulcer and Eradication of H. pylori in a Developing Country: An 18-Month Follow-Up Study," Scand. J. Gastroenterol. vol. 27, 362-
15 66 (1992)). Therefore there is a great need for a vaccine that can be used to treat infection and to prevent future infections.

20 At the present time little is known regarding the role of the mucosal immune systems in the stomach. The distribution of immunoglobulin (Ig) producing cells in the normal gastric antrum indicates that IgA plasma cells make up 80% of the total plasma cell population. In addition, the number of plasma IgA cells present in the gastric antrum is comparable to other mucus membranes. (Brandtzaeg, P. "Role of J Chain and Secretory Component in Receptor-Mediated Glandular and Hepatic Transport of

Immunoglobulins in Man" Scand. J. Immunol., vol. 22, 111-146
(1985); Brandtzaeg, P. et al. "Production and Secretion of
Immunoglobulins in the Gastrointestinal Tract" Ann. Allergy, vol.
59, 21-39 (Nov., 1987)). A number of studies in humans (Wyatt,
5 J.I. et al. "Local Immune Response to Gastritis Campylobacter in
Non-ulcer Dyspepsia" J. Clin. Path., vol. 39, 863-870 (1986)),
and in animal models (Fox, J.G. et al. "Helicobacter mustelae-
Associated Gastritis in Ferrets: An Animal Model of Helicobacter
pylori Gastritis in Humans" Gastroenterology, vol. 99, 352-361
15 (1990); Fox, J.G. et al. "Helicobacter felis Gastritis in
Gnotobiotic Rats: An Animal Model of Helicobacter pylori
Gastritis" Infect. Immun., vol. 59, 785-791 (1991); Fox, J.G. et
al. "Local and Systemic Immune Responses in Murine Helicobacter
felis Active Chronic Gastritis," Infect. & Immun., vol. 61, 2309-
15 (1993)), have demonstrated specific IgG and IgA responses in
serum and in gastric secretions in response to Helicobacter
infection. However, the observation that Helicobacter pylori
infection persists as a chronic infection for years, despite
inducing a local and systemic immune response, is not encouraging
20 the development of immunization strategies.

Lee et al. have reported the ability to infect germ-
free rodents with Helicobacter felis, a bacterium closely related
to Helicobacter pylori, and reproducibly document histologic
gastritis. (Lee, A. et al. "A Small Animal Model of Human

0955739-091801
108T60-6E255660

Helicobacter pylori Active Chronic Gastritis" Gastroenterology, vol. 99, 1315-1323 (1990); Fox, J.G. et al. "Helicobacter felis Gastritis in Gnotobiotic Rats: An Animal Model of Helicobacter pylori Gastritis" Infect. Immun., vol. 59, 785-791 (1991)).

5 Since then, this bacterium-host pairing has been accepted as a good model to study Helicobacter-mediated gastritis and its initiating factors. (Lee, A. et al. "Pathogenicity of Helicobacter pylori: A Perspective" Infect. Immun., vol. 61, 1601-1610 (1993)). Infection of mice with H. felis results in a similar pathologic response to that found in humans infected with H. pylori; both types of infections result in active, chronic gastritis. (Lee et al., Gastroenterology, vol. 99, pp. 1315-1323 (1990)). Researchers have found that Helicobacter felis has the same susceptibility to antimicrobial therapy as Helicobacter pylori, and the H. felis/mouse model has been used to develop new treatments against H. pylori infection. (Dick-Hegedus, E. and Lee, A., "Use of a Mouse Model to Examine Anti-Helicobacter pylori Agents," Scand. J. Gastroenterol., vol. 26, 909-915 (1991); Chen et al., "Immunization Against Gastric Helicobacter Infection in a Mouse/Helicobacter felis Model," Lancet, vol. 339, p.1120 (1992)). Czinn et al. have shown that repetitive oral immunization with a crude lysate of Helicobacter pylori plus cholera toxin adjuvant induces a vigorous gastrointestinal IgA anti-Helicobacter pylori response in mice and ferrets. (Czinn,

10655739-091901
S.J. and Nedrud, J.G. "Oral Immunization Against Helicobacter pylori" Infect. Immun., vol. 59, 2359-2363 (1991)). In addition, Chen et al. and Czinn et al. have recently reported that oral immunization with a crude lysate of H. felis induced protection against H. felis infection in mice. (Chen, et al. "Immunization Against Gastric Helicobacter Infection in a Mouse/Helicobacter felis Model, "(letter) Lancet, vol. 339,1120-1121 (1992); Czinn, S. et al. "Oral Immunization Protects Germ-Free Mice Against Infection from Helicobacter felis" Proceedings of the DDW, American Gastroenterological Association, 1321, A-331 (May 10-13, 1992); Czinn et al., Vaccine, vol. 11, 637-42 (1993)). The exact nature of the antigen(s) responsible for the induction of this protection, however, had not been determined, and no information suggested that the protective antigen(s) of H. felis that induced protection against this pathogen would induce a cross-reactive protection extending to another Helicobacter species.

20 We have demonstrated for the first time that Helicobacter pylori and H.felis shared antigenic determinants by obtaining monoclonal antibodies after oral immunization of mice with either Helicobacter pylori or H. felis sonicates and showing that some of these antibodies, directed against Helicobacter pylori, would crossreact with H. felis and vice versa, (Michetti, P. et al. " Specificity of Mucosal IgA Response in Balb/C Mice Following H. felis or Helicobacter pylori Challenges" Proceedings

of the DDW, American Gastroenterological Association, 1001, A-251
(May 10-13, 1992); Davin, C. et al. "Helicobacter pylori Urease
Elicits Protection Against H. felis Infection in Mice"
Proceedings of the DDW, American Gastroenterological Association,
5 1213, A-304 (May 16-19, 1993)), but the basis for these cross-
reactivities were unknown.

Based on the homology existing between the different
known urease amino acid sequences, it has been proposed that jack
bean urease could be used as a vaccine against Helicobacter
10 pylori. (Pallen, M.J. and Clayton, C.L. "Vaccination Against
Helicobacter pylori Urease" Lancet, vol. 336, 186-7 (1990)).
Nevertheless, despite the homology among the different urease
sequences, cross-reactivity is not the rule. Guo and Liu have
shown years ago that ureases of Proteus mirabilis, Proteus
15 vulgaris and Providencia rettgeri show cross-reactivity to each
other, while ureases of jack bean and Morganella morganii are
immunologically distinct from the three former ureases. (Guo, M.
and Liu, P.V. "Serological Specificities of Ureases of Proteus
Species" J. Gen. Microbiol, vol. 136, 1995-2000 (1965)). So,
20 even if an antigenic cross-reactivity of Helicobacter pylori
urease with other Helicobacter ureases was a reasonable postu-
late, no data existed demonstrating that this was really the case
until we showed that some H.felis monoclonal antibodies cross-
reacted with Helicobacter pylori urease. (Davin, C. et al.

5 "Helicobacter pylori Urease Elicits Protection Against H. felis
Infection in Mice" Proceedings of the DDW, American
Gastroenterological Association 1213, A-304 (May 16-19, 1993)).
J. Pappo has further demonstrated that mice which have been
infected by H. felis produce antibodies which crossreact with
Helicobacter pylori urease but not jack bean urease (J. Pappo,
unpublished data, 1993). The fact that jack bean urease does not
fall in the same immunological category than Helicobacter urease
suggests that jack bean urease may not be useful for immunization
against Helicobacter infections, the way it was done for enteric
bacteria. (Pimentel, J.L. and Cook, M.E. "Improved Growth in the
Progeny of Hens Immunized with Jackbean Urease" Poultry Sci.,
vol. 64, 434-439 (1988)). Furthermore, attempts to immunize mice
against H. felis infection by oral or intraperitoneal delivery of
jack bean urease resulted in the production of antibodies against
jack bean urease, but failed to protect the mice from infection.
(Chen, M. et al. "Failure of Immunization Against Helicobacter
Using Jack Bean Urease," Acta Gastroenterol. Belg., vol. 56, 94
(1993)).

20 The use of an antigen that is the reaction product of
urease and glutaraldehyde is described in U.S. Patent No.
4,837,017, "Urease Antigen Product and Process," issued June 6,
1989, to LeVeen et al. The patent describes the use of the
antigen to reduce ammonia toxicity caused by urea splitting

organisms. LeVeen et al. disclose the injection of glutaral-
dehyde treated jack bean urease into the bloodstream. The LeVeen
patent does not disclose the administration of the urease antigen
to the mucosal surface of a mammal in order to stimulate antibody
5 production by the local immune system. Furthermore, there is no
evidence in the specification that the injection of a jack bean
urease antigen could prevent Helicobacter infection or be used to
treat gastroduodenal infection by Helicobacter.

Eaton et al. have shown that mutant H. pylori cultures
with weak urease activity are unable to infect gnotobiotic
piglets. (Eaton et al., "Essential Role of Urease in the
Pathogenesis of Gastritis Induced by Helicobacter pylori in
Gnotobiotic Piglets," Gastroenterology, vol. 98, A654 (1990)).
Eaton does not describe the use of a urease antigen as a vaccine
to prevent Helicobacter infection or as a method of treating
15 Helicobacter infection.

The use of Helicobacter pylori urease, or of related
ureases, as a vaccine against Helicobacter pylori infection has
previously been proposed by A. Labigne, and incorporated among
20 the claims of a patent filed on October 6th, 1988 by Pasteur
Institute, Paris, France. (Labigne, A. "Sequences of Nucleotides
Coding for a Protein Having an Urease Activity". EPO patent
application # EPO 367 644 A1, 1989. International Publication
WO 90/04030, 1990). The specification of this document con-

5 tains, however, no evidence of vaccination of any mammal against any Helicobacter infection with urease. This part of the Pasteur Institute patent, therefore, has not been reduced to practice, and the related claims (claims 27 and 28, page 16) should not be considered as valid. Furthermore, the claims of this document relate to a protein presenting a urease activity, and it will be understood from the experiments described below that enzymatic activity of the urease-based vaccine is not required to induce protection after oral immunization.

10
15
20 Moreover, while sequence homology with other bacterial ureases might support the use of urease as a vaccine candidate against Helicobacter pylori infection, the current knowledge of human Helicobacter pylori infection would certainly not. First, despite the fact that infected individuals often mount a strong antibody response to urease, the anti-urease immune response does not result in clearance or control of the infection. Second, Helicobacter pylori is able to transport urease out of the cell and to shed it from its surface, (Evans, D.J. et al. "Urease-Associated Heat Shock Protein of Helicobacter pylori" Infect. Immun., vol. 60, 2125-2127 (1992), Ferrero, R.L. and Lee, A. "The Importance of Urease in Acid Protection for the Gastric-Colonizing Bacteria Helicobacter pylori and Helicobacter felis sp. nov." Microb. Ecol. Health Dis., vol. 4, 121-134 (1991)), thus urease may not represent an appropriate target for the

development of a protective mucosal immune response. Indeed, mucosal immune protection is thought to be mainly mediated by secretory IgA, the agglutinating activity of which would be impaired when the recognized antigen can be shed by the target pathogen. Third, urease appears to be toxic for epithelial cells in culture, and has been suspected to play a role in mucus degradation and in peptic ulceration in vivo (Mégraud, F. et al., "Further Evidence of the Toxic Effect of Ammonia Produced by Helicobacter pylori Urease on Human Epithelial Cells," Infect. & Immun., vol. 60, 1858-63 (1992); Murakami, M. et al., "Gastric Ammonia has a Potent Ulcerogenic Action on the Rat Stomach," Gastroenterology 1993, vol. 105, 1710-15), thus its use as antigen may be toxic.

Nevertheless, we reasoned that this antigen could be a potentially efficient vaccine if:

- first, we would deliver it orally at a sufficiently high dose to elicit a stronger immune response than the naturally occurring one

- second, the amount of antibodies produced would be high enough to bind all the urease, shed or not shed

- third, we would use subunits of urease or a molecular species that was non toxic.

Another aspect of the invention describes the use of antibodies directed against urease to prevent and to treat

Helicobacter infection. European Patent Application No. 91310049.1, filed by Kunio Ando on October 31, 1991, claiming priority on Japanese Patent Application No. 296609/90 filed November 1, 1990, titled "A Method for Producing a new Medicine for Both Treating and Preventing Peptic Ulcer Diseases and Gastritis and Thus Formulated Medicines," describes the oral administration of polyclonal antibodies derived from bovine colostrum and bovine serum to patients with active chronic gastritis type B and to patients with duodenal ulcer. The Ando application describes the use of an antibody preparation directed against many antigens, including Helicobacter pylori, and does not disclose the use of an antibody directed against urease to treat or prevent Helicobacter pylori infection. The use of antibodies to treat gastric disease in gnotobiotic piglets was described in United States Patents No. 5,258,178 and 5,260,057, issued to Cordle and Schaller and titled "Method and Product for the Treatment of Gastric Disease." The Cordle and Schaller patents describe the use of an antibody preparation that does not solely contain antibodies directed against Helicobacter pylori, and does not disclose the use of an antibody directed against urease to treat or prevent Helicobacter pylori infection. Nagata et al. describe the preparation of a monoclonal antibody directed against Helicobacter pylori that inhibits urease activity. (Nagata, K., et al., "Monoclonal Antibodies Against the Native

Urease of Helicobacter pylori: Synergistic Inhibition of Urease Activity by Monoclonal Antibody Combinations," Infect. and Immun., Vol. 60, 4826 (1992)). Nagata et al. do not describe the use of monoclonal antibodies directed against urease to prevent or to treat Helicobacter pylori infection.

Very few examples of therapeutic vaccines are available in the literature. Most of them are related to parenteral immunizations aimed to stimulate the host's immune system against malignant tumors, to modulate the immune system in autoimmune diseases such as rheumatoid arthritis or as desensitization in allergy states. Therapeutic vaccination procedures against different infections were also performed, most of them via a parenteral route of immunization. They included immunizations against leprosy in humans (Zaheer SA et al. "Combined Multidrug and Mycobacterium w Vaccine Therapy in Patients with Multibacillary Leprosy" J. Infect Dis., vol. 167, 401-410 (1993), Mukherjee A. et al., "Histopathological Monitoring of an Immunotherapeutic Trial with Mycobacterium w." Int. J Lepr. Other Mycobact. Dis., vol. 60, 28-35 (1992)), in complementation of antibiotic therapy, vaccination against Phythiosis insidiori, a mycological infection, in horses (Mendoza L, et al., "Evaluation of Two Vaccines for the Treatment of Pythiosis insidios in Horses" Mycopathologia, vol. 119, 89-95 (1992)), an uncontrolled study on the use of an autovaccine in chronic osteomyelitis

(Sologub VV. "Experience in Using an Autovaccine in Treating Patients with Chronic Osteomyelitis" Vrach, Delo, 122-125 (1992)) and systemic immunization against Campylobacter fetus infection of female cattle (Schurig, G.G.D., et al., "Bovine Venereal Vibriosis: Cure of Genital Infection in Females by Systemic Immunization," Infect. & Immun., Vol. 11, 245-51 (1975)). To date, only one oral immunotherapy study aimed at stimulating the mucosal immune system in order to treat (and to prevent recurrence of) a mucosal infection has been performed, for urinary tract infection (Schulman CC, et al. "Oral Immunotherapy of Recurrent Urinary Tract Infections: A Double-Blind Placebo-Controlled Multicenter Study" J Urol., vol. 150, 917-921 (1993)). In that study, Schulman et al. used a lysate of selected E. coli strains, together with an concomitant treatment of antibiotics, chemotherapeutics or urinary tract disinfectants to treat the acute infection at entry in the study. Therefore, no study has demonstrated so far the effectiveness of a therapeutic vaccine, used as a monotherapy, administered to the mucosal immune system, against a bacterial disease.

The novelty of a therapeutic vaccine against Helicobacter infection also comes from the observation that H. pylori persists as a chronic infection in the gastric cavity for years, despite inducing a vigorous local and systemic immune response. This observation was conceptually already an obstacle

to the development of a prophylactic vaccine against Helicobacter infection, but was even more an obstacle to the development of a therapeutic immunization.

In summary, there remains a need for effective treatment and prevention of Helicobacter pylori-induced gastric infection in humans. Recent data suggested the possibility to generate a vaccine against this infection, but have not provided a clear identification of defined antigen(s), common to all strains of Helicobacter pylori, that could be incorporated into a safe and effective vaccine.

In this invention, we have identified the urease antigen of Helicobacter pylori as a candidate vaccine and demonstrated its efficacy in an animal model. We have also demonstrated the use of the Helicobacter pylori urease antigen for the treatment and eradication of Helicobacter infection. We have further demonstrated that the B subunit of urease alone (ure B) is effective as a vaccine useful for the prevention of and treatment of Helicobacter infection. These results were unexpected in the light of the natural history of Helicobacter infections.

SUMMARY OF THE INVENTION

We have determined that it is useful to immunize animals with Helicobacter urease peptides for both prophylactic and therapeutic treatment. Immunization of animals with Helicobacter urease peptides prevents infection by Helicobacter and eradicates infection in previously infected animals. This method, and the vaccine compositions, are useful for the prevention and treatment of gastroduodenal disease associated with Helicobacter infection.

We have discovered that immunity can be induced in mammals susceptible to gastrointestinal Helicobacter infection by exploiting urease epitopes displayed on or about the surface of Helicobacter organisms and using them as a vaccine target. The immunity can be induced by immunization with native urease, but can also be induced with recombinant urease subunit, produced as an enzymatically inactive, therefore non-toxic form. The invention provides a method of inducing immunity to Helicobacter infection by administering to a mucosal surface of a mammal a polyaminoacid preparation, i.e. a mixture of peptides and/or proteins, together with an appropriate adjuvant. This polyaminoacid preparation presents a plurality of epitopes characteristic of and exhibited by a urease enzyme endogenous to the infecting Helicobacter organism. The administration of the polyaminoacid preparation may be performed by the oral route.

The active ingredient of the preparation may comprise natural or biosynthetic epitopes and may take various forms. A non exhaustive list of possible preparations includes purified, naturally occurring or recombinantly produced urease preparations of bacterial or other origin, digests of urease, fusion proteins comprising urease epitopes, truncated forms of urease enzyme, or peptides homologous with the amino acid sequence of urease. Since development of immunity depends on induction of humoral and/or cellular immune responses which bind to the infecting Helicobacter organism, preferred preparations are those which most closely duplicate the epitopes of the urease endogenous to the infecting organism. For example, preparations displaying the epitopes of urease of Helicobacter pylori are preferred for administration in humans susceptible to Helicobacter pylori, and preparations displaying the epitopes of urease of H. felis are preferred for the administration in humans susceptible to H. felis. However, in accordance with an important aspect of the invention, it has been discovered that urease from a heterologous species may be used. For example, we have shown that H. felis infection in mice can be prevented by administration of urease from Helicobacter pylori. Thus, H. pylori urease can be used to protect against H. pylori as well as H. felis. H. felis is an occasional cause of human infection and disease. (Wegman, W. et al., Schweig. Med. Wochenschr. vol. 121, 245-54 (1991)).

According to a first embodiment of the present invention, a method is provided of eliciting in a mammalian host a protective immune response to Helicobacter infection.

According to a second embodiment of the present invention, a method is provided of treating a mammalian host that is infected with Helicobacter.

Thus, in a first aspect, the present invention provides a method of eliciting in a mammalian host a protective immune response to Helicobacter infection. The method comprises the step of administering to a mucosal surface of the mammal, including humans, an immunologically effective amount of a urease antigen, preferably Helicobacter pylori urease, capable of eliciting such a protective immune response. The term "comprising" is used herein as it is recognized in the art.

According to one aspect of the invention, there is provided a method of eliciting in a mammalian host a protective immune response to Helicobacter infection wherein an immunologically effective amount of a urease antigen capable of eliciting such a protective immune response, preferably Helicobacter pylori urease or Helicobacter pylori urease B subunit, is administered to a mucosal surface of the host.

In a second aspect, the present invention provides a method of eliciting in a mammalian host a protective immune response to Helicobacter infection. The method comprises the step

of administering to a mucosal surface of the mammal, including humans, an immunologically effective amount of recombinant, enzymatically inactive, urease B subunit as antigen, preferably recombinant Helicobacter pylori urease B subunit, capable of eliciting such a protective immune response.

The invention also includes within its scope the treatment or prophylaxis of mammals, including humans, for Helicobacter infection, wherein an immunologically effective amount of a urease, or its subunits, capable of eliciting a protective immune response to Helicobacter infection, is administered to a mucosal surface of a patient. Preferably, the urease is Helicobacter pylori urease or Helicobacter pylori urease B subunit, and the urease is preferably administered in particulate form in association with a hydroxylated calcium phosphate, for example hydroxyapatite. Moreover, it is preferred to administer the Helicobacter pylori urease in association with a mucosal adjuvant, the B subunit of cholera toxin, muramyl dipeptide or other such adjuvants.

According to another aspect of the present invention, there is provided a vaccine composition suitable for prevention of Helicobacter infection, comprising an effective amount of a urease antigen, preferably Helicobacter pylori urease or Helicobacter pylori urease B subunit, or recombinant Helicobacter pylori urease subunits, capable of eliciting in a host a

protective immune response to Helicobacter infection, in association with a pharmaceutically acceptable carrier or diluent. Suitable carriers and diluents will be recognized by those skilled in the art and can be found in, for example

5 Remington's Pharmaceutical Sciences (18th ed., 1990).

10
15
20
The vaccines of the invention are administered in amounts readily determined by persons of ordinary skill in this art. Thus, for adults a suitable dosage will be in the range of 10 μ g to 100 milligrams, for example 50 μ g to 50mg. A suitable dosage for adults will also be in the range of 5 μ g to 500mg. Similar dosage ranges will be applicable for children. Carrier systems in humans may include enteric release capsules protecting the antigen from the acidic environment of the stomach, and including urease antigen in a insoluble form as fusion proteins. The vaccine can be administered as a primary prophylactic agent in adults or in children, as a secondary prevention, after successful eradication of Helicobacter pylori in an infected host, or as a therapeutic agent in the aim to induce an immune response in the host susceptible to contribute to the eradication of Helicobacter pylori.

As noted above, a suitable mucosal adjuvant is cholera toxin. Others which may be used are non-toxic derivatives of cholera toxin, including its B subunit, and/or conjugates or genetically engineered fusions of the urease antigen plus cholera

toxin or its B subunit. Other suitable delivery methods include biodegradable microcapsules or immuno-stimulating complexes (ISCOMs) or liposomes, genetically engineered attenuated live vectors such as viruses or bacteria, and recombinant (chimeric) virus-like particles, e.g. bluetongue. The amount of mucosal adjuvant employed depends on the type of mucosal adjuvant used. For example, when the mucosal adjuvant is cholera toxin, it is suitably used in an amount of 5 μ g to 50 μ g, for example 10 μ g to 35 μ g. When used in the form of microcapsules, the amount used will depend on the amount employed in the matrix of the microcapsules to achieve the desired dosage. The determination of this amount is within the skill of a person of ordinary skill in this art.

Suitable carriers for the vaccines of the invention are enteric coated capsules and polylactide-glycolide microspheres. Suitable diluents are 0.2N NaHCO₃ and/or saline.

Particulate hydroxylated calcium phosphate (HCP) is especially useful as a carrier for the Helicobacter pylori urease to be applied to mucosal surfaces. It is believed that the Helicobacter pylori urease-hydroxylated calcium phosphate conjugate is transported across epithelium where it raises a poly Ig immune response. Preferably, the hydroxylated calcium phosphate is in the form of microparticles suitable for transport across the epithelium, particularly by cells specialized for this

purpose (M cells). A preferred form of hydroxylated calcium phosphate is hydroxyapatite, a commercially available crystalline hydroxylated calcium phosphate $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$.

Commercially available hydroxyapatite generally consists of slab-like crystals that are chemically and physically analogous to inorganic hydroxyapatite in normal bone tissue. Ingestion of hydroxyapatite should therefore be safe, as evidenced by the existence of nutritional calcium/phosphorus supplements derived from ground bone, which are designed to be ingested. Commercially-high resolution hydroxyapatite (from CalBiochem) consists of crystals varying widely in size. Crystals over $1\mu\text{m}$ in length are unlikely to be taken up by M cells. Therefore, for use in the invention, commercial hydroxyapatite crystals are broken into small, relatively uniform crystalline fragments such as by sonication. Preferably, a substantial proportion of the hydroxyapatite is present as fragments of about $0.01\text{-}1.0\mu\text{m}$. Fragmentation may be measured either by electron microscopy or light scattering, using standard techniques.

Preferred modes of administration of the Helicobacter pylori urease antigen are orally, nasally, rectally or ocularly. Oral administration can provide delivery to other G.I. (gastrointestinal) mucosa including the intestinal mucosa.

The vaccines of the present invention may be administered to a mucosal surface in the form of an aerosol,

suspension, capsule and/or suppository. The method of administration will be readily apparent to a person of ordinary skill in this art and can be found in, for example, Remington's Pharmaceutical Sciences (18 ed., 1990).

5 According to a further aspect of the present invention, there is provided a method of imparting to a mammalian host passive protection to Helicobacter infection, comprising administering to a mucosal surface of the host an immunologically effective amount of a urease specific antibody produced in a host immunized with a urease, preferably Helicobacter pylori urease or Helicobacter pylori urease B subunit, capable of eliciting a protective immune response to Helicobacter infection.

10 The present invention further includes the passive immunization of mammals, including humans, against Helicobacter infection. This is achieved by administering to a mucosal surface of the patient an effective amount of a urease specific antibody, preferably, an effective amount of a Helicobacter pylori urease specific IgA monoclonal antibody.

15 Since the urease of Helicobacter pylori is shown to
20 represent the antigen involved in inducing protective immunity, a further aspect of the invention is the use of Helicobacter pylori urease as a diagnostic reagent to measure the immune response of persons who have received a vaccine based on urease or to determine whether an individual is immune or susceptible (and thus in

0055739.091301
1055739.091301

need of vaccination). The present invention also includes the use of urease and urease-specific antibodies to construct assays and kits for diagnosis of Helicobacter immunity, assessment of Helicobacter susceptibility, and definition of immune responses to vaccines.

In a third aspect, the invention provides a method of treating gastroduodenal disease in a mammal. This method comprises the step of administering a therapeutically effective amount of a composition comprising Helicobacter urease peptides. The gastroduodenal diseases included within the scope of the invention include, but are not limited to, gastritis, peptic ulcer disease, including both gastric and duodenal ulcers, gastric cancer, chronic dyspepsia with severe erosive gastroduodenitis, refractory ulcer dyspepsia, intestinal metaplasia, low grade MALT lymphoma, Helicobacter infection, Helicobacter pylori infection and Helicobacter felis infection. The term "urease peptides" refers to, but is not limited to, any urease or subunit or urease, either naturally occurring or obtained by recombinant DNA techniques, as well as a digested fragment or peptide thereof, fusion proteins comprising the whole urease, subunits, or fragments thereof, or truncated urease constructs. Also included within the term "urease peptides," are proteins or peptides that display epitopes sufficiently homologous to epitopes displayed by Helicobacter urease such that antibodies

that recognize epitopes displayed by Helicobacter urease will recognize epitopes displayed by said peptides or proteins.

In a more particular related aspect, the invention provides a method of treating gastroduodenal disease caused by Helicobacter infection. The Helicobacter infection may be, but is not limited to, Helicobacter pylori or H. felis infection.

More particularly, the invention provides a method of treating gastroduodenal disease caused by Helicobacter infection whereby the composition comprising Helicobacter urease peptides is administered to a mucosal surface. Without limiting the type of mucosal surface used for administration, the mucosal surface may be oral, nasal, rectal, or ocular.

The invention can also feature the administration of a composition comprising Helicobacter urease peptides in association with a mucosal adjuvant. The mucosal adjuvant may be selected from, but is not limited to, cholera toxin, procholera toxin, cholera toxin B subunit, fungal polysaccharides including, but not limited to, schizophyllan, muramyl dipeptide, muramyl dipeptide derivatives, phorbol esters, microspheres, non-Helicobacter pylori bacterial lysates, labile toxin of Escherichia coli, block polymers, saponins, and ISCOMs. Other mucosal adjuvants will be recognized to those in the art and can be found in, for example, Azuma, I., "Synthetic Immunoadjuvants: Application to Non-Specific Host Stimulation and Potentiation of

0955739-091901
104160-6375560

Vaccine Immunogenicity" Vaccine, vol. 10, 1000 (1992); Pockley,
A.G. & Montgomery, P.C., "In vivo Adjuvant Effect of Interleukins
5 and 6 on Rat Tear IgA Antibody Responses" Immunology, vol. 73,
19-23 (1991); Adam, A. & Lederer, E. "Muramyl peptides as
5 Immunomodulators" ISI ATLAS OF SCIENCE 205 (1988); Clements,
J.D., et al. "Adjuvant Activity of Escherichia coli Heat-labile
Enterotoxin and Effect on the Induction of Oral Tolerance in Mice
to Unrelated Protein Antigens" Vaccine, vol. 6, 269 (1988); Ben
Ahmeida, E.T.S., et al. "Immunopotential of Local and Systemic
Humoral Immune Responses by ISCOMs, Liposomes and FCA: Role in
10 Protection Against Influenza A in Mice" Vaccine, vol. 11, 1302
(1993); and Gupta, R.K. et al. "Adjuvants -- A Balance Between
Toxicity and Adjuvanticity" Vaccine, vol. 11, 290-308 (1993).

The mucosal adjuvant may also be genetically or
chemically linked to the urease peptides. Examples of this type
of fusion peptide are known to those skilled in the art and can
also be found in Czerkinsky et al., "Oral Administration of a
Streptococcal Antigen Coupled to Cholera Toxin B Subunit Evokes
Strong Antibody Responses in Salivary Glands and Extramucosal
15 Tissues" Infect. Immun., vol. 57, 1072-77 (1989); Nashar et al.,
"Current Progress in the Development of the B Subunits of Cholera
Toxin and Escherichia Coli Heat-Labile Enterotoxin as Carriers
for the Oral Delivery of Heterologous Antigens and Epitopes"
20 Vaccine, vol. 11, 235-40 (1993); and Dertzbaugh and Elson,

"Comparative Effectiveness of the Cholera Toxin B Subunit and Alkaline Phosphatase as Carriers for Oral Vaccines," Infect. Immun., vol. 61, 48-55 (1993). For example, the urease B subunit could be expressed as a chimeric protein that is genetically
5 linked to the cholera toxin B subunit through the use of a DNA expression vector containing the ure B nucleotide sequence linked to the cholera toxin B subunit nucleotide sequence.

In another related aspect, the method involves the administration of a composition comprising Helicobacter urease peptides where such composition is delivered in particulate form. The composition may be delivered in particulate form through association with a carrier. The carrier may be a hydroxylated calcium phosphate, for example, hydroxyapatite. The term
10 "hydroxyapatite" refers to, but is not limited to, its meaning as recognized by those skilled in the art to mean a tribasic calcium phosphate, also known as hydroxylated calcium phosphate or calcium hydroxide phosphate. This is only an example, and is not
15 meant to be limiting as to the type of carrier that may be used.

In another related aspect, the administered dosage of
20 the composition comprising Helicobacter urease peptides may range from 100 μ g to 1g, for example, 0.14 mg to 14.4 mg per kg of body weight. Those of skill in the art will recognize that the optimal dose may be more or less depending upon the patient's body weight, disease, the route of administration, and other

factors. The dosage level is readily determinable by standard methods. The number of doses will depend upon the disease, the formulation, and efficacy data from clinical trials. For example, the dosage may be administered over 3 to 8 doses for a primary immunization schedule over 1 month. This course of treatment is an example and is not meant to be limiting.

In one related aspect, the method involves the administration of a composition comprising Helicobacter urease peptides wherein the composition is administered in association with a microsphere carrier. Such microsphere carrier may be, for example, but is not limited to, a polylactide-coglycode biodegradable microsphere carrier.

In another related aspect, the method involves the administration of a composition comprising the Helicobacter urease peptides wherein such composition comprises a recombinant live vector which expresses a Helicobacter urease peptide. Those skilled in the art will recognize that such live vector may be, for example, a bacterial or a viral vector. For example, without any limitation, the live vector may be selected from the group consisting of Salmonella typhimurium, Salmonella typhi, Shigella, Bacillus, Lactobacillus, BCG, Escherichia coli, Vibrio cholerae, Campylobacter, yeast, Herpes virus, Adenovirus, Poliovirus, Vaccinia, and Avipox. In addition, a carrier system which expresses a Helicobacter urease peptide, such as Bluetongue

virus-like particles, Rotavirus-like particles, and Ty particles, may be used to deliver the urease peptide. In a preferred aspect, the live vector or the carrier system may be administered to a mucosal surface.

5 A preferred embodiment of the present invention comprises a method of treating a human infected with Helicobacter pylori, comprising orally administering a therapeutically effective amount of a composition comprising the ure B subunit of Helicobacter pylori urease, in association with a mucosal adjuvant selected from the group consisting of cholera toxin, procholera genoid, cholera toxin B subunit, fungal polysaccharides, including schizophyllan, muramyl dipeptide, muramyl dipeptide derivatives, phorbol esters, liposomes, microspheres, non-Helicobacter pylori bacterial lysates, labile toxin of Escherichia coli, block polymers, saponins, and ISCOMs, the composition being administered in a particulate form in association with hydroxyapatite. The composition may also be administered with concurrent oral administration of a chewable 1.0g NaHCO₃ tablet. For the purposes of this invention, the term "in association with" includes any type of association, including but not limited to, a chemical or genetic association, such as that present in a fusion protein.

In a fourth aspect, the invention provides a method of treating gastroduodenal disease in a mammal, comprising adminis-

tering a therapeutically effective amount of a composition comprising an antibody that recognizes Helicobacter urease. The gastroduodenal diseases included within the scope of the invention are referenced above. These methods use the above-referenced composition comprising Helicobacter urease peptides to elicit an antibody response in a mammal. The antibodies produced by the immunized mammal are isolated and administered to the subject mammal. The preparation of antibodies that recognize a given antigen, such as the above-referenced composition, is known to those skilled in the art. For example, polyclonal and monoclonal antibodies can be prepared following the disclosure in Harlow, E. & Lane, D., Antibodies: A Laboratory Manual (1988).

In a more particular related aspect, the administered antibody is a monoclonal antibody. The preparation of monoclonal antibodies is known to those skilled in the art.

More particularly, the administered antibody is an IgA antibody, either a polyclonal or monoclonal IgA antibody. The preparation of monoclonal IgA antibodies is known to those skilled in the art and may be found in, for example, Winner, L., et al., "New Model for Analysis of Mucosal Immunity: Intestinal Secretion of Specific Monoclonal Immunoglobulin A from Hybridoma Tumors Protects Against Vibrio Cholerae Infection" Infect. and Immun., vol. 59, 977-982 (1991); and Weltzin, R., et al., "Binding and Transepithelial Transport of Immunoglobulins by

0955739-091301
103160-5E255660

Intestinal M Cells: Demonstration Using Monoclonal IgA
Antibodies Against Enteric Viral Proteins" J. Cell Biol., vol.
108, 1673-1685 (1989).

5 The term "antibody" as used for the purposes of this
invention includes, but is not limited to, polyclonal antibodies,
monoclonal antibodies, and anti-idiotypic antibodies. The
antibodies may be naturally derived from any animal, synthesized
in bacteria or another non-animal source, chemically synthesized,
or genetically synthesized.

10 In a preferred aspect of the invention, a method is
provided of treating a human infected with Helicobacter pylori,
comprising administering a therapeutically effective amount of a
composition comprising an IgA monoclonal antibody that recognizes
the ure B subunit of Helicobacter pylori urease.

15 In a fifth aspect, the invention provides a composition
useful in the therapeutic treatment of gastroduodenal disease,
comprising Helicobacter urease peptides. The gastroduodenal
diseases included within the scope of the invention, as well as
the Helicobacter urease peptides and mucosal adjuvants included
20 within the scope of the invention, are referenced above.

In a related aspect, the composition comprises
Helicobacter urease peptides and a mucosal adjuvant.

In another related aspect, the composition is in
particulate form. The composition may exist in particulate form

through association with a carrier. The carrier may be, for example, hydroxyapatite, as referenced above. This is only an example, and is not meant to be limiting as to the type of carrier that may be used.

5 More particularly, the composition is present in particulate form, in a liquid suspension.

10 In a related aspect, the composition comprises Helicobacter urease peptides in association with a microsphere carrier. Such microsphere carrier may be, for example, but is not limited to, a polylactide-coglycolide biodegradable microsphere. The urease peptides may encapsulated in the biodegradable microspheres. Polylactide-coglycolide microspheres slowly hydrolyze in the presence of water and becomes water-soluble, thus delivering the peptides that are incorporated in the microsphere.

15 In another related aspect, the composition comprises Helicobacter urease peptides wherein the composition comprises a recombinant live vector which expresses a Helicobacter urease peptide. Those skilled in the art will recognize that such live
20 vector may be, for example, a bacterial or a viral vector. for example, without any limitation, the live vector may be selected from the group consisting of Salmonella typhimurium, Salmonella typhi, Shigella, Bacillus, Lactobacillus, BCG, Escherichia coli, Vibrio cholerae, Campylobacter, yeast, Herpes virus, Adenovirus,

Poliovirus, Vaccinia, and Avipox. In addition, a carrier system which expresses a Helicobacter urease peptide, such as Bluetongue virus-like particles, Rotavirus-like particles, and Ty particles, may be used to deliver the urease peptide.

5 In a preferred embodiment, a composition is provided that is useful in the therapeutic treatment of Helicobacter pylori infection of a human, comprising the ure B subunit of Helicobacter pylori urease, a mucosal adjuvant selected from a group consisting of cholera toxin, procholera toxin, cholera toxin B subunit, fungal polysaccharides, including schizophyllan, muramyl dipeptide, muramyl dipeptide derivatives, phorbol esters, liposomes, microspheres, non-Helicobacter pylori bacterial lysates, labile toxin of Escherichia coli, block polymers, saponins, and ISCOMs, and further comprising hydroxyapatite, wherein the composition is in a particulate form in a liquid suspension. The term "in association with" is defined as referenced above.

10 In a sixth aspect, a composition is provided that is useful in the therapeutic treatment of gastroduodenal disease,
20 comprising an antibody that recognizes Helicobacter urease. The gastroduodenal disease included within the scope of the invention, as well as the antibodies included within the scope of the invention, are referenced above.

In a preferred embodiment of the invention, a composition is provided that is useful in the therapeutic treatment of gastroduodenal disease, comprising an IgA monoclonal antibody that recognizes the ure B subunit of Helicobacter pylori urease.

5 While not being bound by any theory, the present inventors believe that administration of the urease antigen, or B subunit thereof, to a mucosal surface stimulates the common mucosal immune system and perhaps local sites in the gastric mucosa inducing an immune response, including the appearance of Helicobacter pylori specific IgA antibodies in the gastric secretions, which prevent Helicobacter infection. Because of this belief, the terms "immunization" and "vaccine" are used herein by their common meanings as recognized by those skilled in the art and also to indicate methods and compositions used for the treatment of Helicobacter infection, without limitation to these meanings, and without being bound to any theory regarding the mechanism of treatment.

10
15
20 Since it is a routine matter to conduct pre-clinical trials of candidate vaccines for human use in animal models, it is believed that the methodology of the present invention is effective in humans, especially in the prevention and treatment of peptic ulcers, gastritis, gastric malignancies and other conditions arising as a result of the presence of Helicobacter pylori and/or H. felis.

Based on the dosage and course of treatment that successfully eradicated infection in the mouse model, a preferred range of dosage would be 100 μ g to 10 g of Helicobacter urease peptides. Those skilled in the art will recognize the appropriate dosage level to test from research reported for other oral vaccines such as, for example, the research performed with Escherichia coli lysate (6 mg dose daily up to a total of 540 mg) and with an enterotoxigenic E. coli purified antigen (4 doses of 1 mg). (Schulman et al, "Oral Immunotherapy of Recurrent Urinary Tract Infections: A Double-Blind Placebo-Controlled Multicenter Study," J. Urol., vol. 150, 917-921 (1993); Boedeker et al., "Safety, Immunogenicity and Efficacy in Human Volunteers of Biodegradable, Biocompatible Microspheres Containing Colonization Factor Antigen/II (CFA/II) as an Enteral Vaccine Against Enterotoxigenic E. coli (ETEC)" American Gastroenterological Assoc., vol. 999, A-222 (1993)). Without intending any limitation as to the course of treatment, the treatment could be administered over 3 to 8 doses for a primary immunization schedule over 1 month. (Boedeker, American Gastroenterological Assoc., vol. 888, A-222 (1993)).

A recommended method of immunization is administration of the Helicobacter urease peptide composition in the form of a liquid alone or suspension containing Na_2HCO_3 or similar material to temporarily neutralize gastric acid. The neutralizing

material may also be delivered separately at the time the peptide composition is administered, such as in the form of a chewable Na_2HCO_3 tablet. Alternatively, the composition can be administered in the form of enteric-coated capsules. These methods will likely avoid the problem of degradation of the urease composition during its passage in the upper gastrointestinal tract. These methods are reviewed in, for example, Levine & Norlega, "Vaccines to Prevent Enteric Infections," *Ballieres Clin. Gastro.*, vol. 7, 501-517 (1993).

Although a high dose of cholera toxin is not preferred as a mucosal adjuvant for human use, other mucosal adjuvants recognized by those skilled in the art, and as referenced above, will be useful for treatment of humans.

It will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention will now be further described with reference to the accompanying drawings, in which Figures 1 through 8 are graphical representations of the results set forth in Tables 1 through 9.

Figure 1 is a graphic representation of the results from Table 1 of the tests for antibodies in serum (IgG) and intestinal secretion (IgA) in mice that were not protected after immunization with urease.

5 Figure 2 is a graphic representation of the results from Table 1 of tests for antibodies in serum (IgG) and intestinal secretion (IgA) in mice that were protected after immunization with urease.

10 Figure 3 is a graphic representation of the results from Table 1 of tests for antibodies in serum (IgG) and intestinal secretion (IgA) in mice that were not protected after immunization with Helicobacter pylori sonicate.

15 Figure 4 is a graphic representation of the results from Table 1 of tests for antibodies in serum (IgG) and intestinal secretion (IgA) in mice that were protected after immunization with Helicobacter pylori sonicate.

Figure 5 is a graphic representation of the results set out in Table 2 comparing the level of protection obtained with Helicobacter pylori urease as compared to that obtained with Helicobacter pylori sonicate and with cholera toxin.

09055739.091801
103T60" 5255660

Figure 6 is a graphic representation of the results set out in Tables 5 and 6 which measures urease activity in mice that were challenged after oral immunization with recombinant urease A and B subunits.

5 Figure 7 is a graphic representation of the results set out in Table 8 which measures gastric tissue urease activity as a reflection of Helicobacter infection in mice that were subsequently treated with the Helicobacter pylori ure B subunit, cholera toxin and hydroxyapatite only (sham immunized), or untreated.

10
15 Figure 8 is a graphic representation of the results set out in Table 9 which measures gastric tissue urease activity as a reflection of Helicobacter infection in mice that were subsequently treated with the Helicobacter pylori ure B subunit, or cholera toxin and hydroxyapatite only (sham immunized). For Figure 8A, the mice were sacrificed and urease assays performed 2.5 weeks after the last immunization. For Figure 8B, the mice were sacrificed and urease assays performed 8 weeks after the last immunization.

20 DETAILED DESCRIPTION OF THE INVENTION

10655739.091901
10655739.091901

The present inventors have discovered that oral administration to mice of polyaminoacid preparations exhibiting the epitopes of Helicobacter pylori urease gives rise to a protective immunological response against H. felis in mice, an animal model of generally-accepted value for the study of the immune response to Helicobacter infection (Lee, A. et al. "A Small Animal Model of Human Helicobacter pylori Active Chronic Gastritis" Gastroenterology, vol. 99, 1315-1323 (1990)), and a recognized pathogen causing gastritis in humans (Wegman, W. et al., Schweig. Med. Wochenschr., vol. 121, 245-54 (1991)). The effect of the protective immune response is that immunized animals, when challenged with pathogen, have a greatly reduced incidence of infection, in comparison to non-immunized animals. Furthermore, the inventors have discovered that oral immunization in mice using Helicobacter pylori urease B subunit, produced as an enzymatically-inactive recombinant protein, gives rise to a protective immunological response in mice against H. felis. The effect of the protective immune response is that immunized animals, when challenged with pathogen, have also a greatly reduced incidence of infection, in comparison to non-immunized animals which do become infected.

The present inventors have discovered that oral administration of Helicobacter urease peptides to mice infected with H. felis results in the clearance of the infection. This

result indicates that the oral administration of Helicobacter urease peptides is an effective therapy for the treatment of Helicobacter infection in mammals. The oral administration of Helicobacter pylori urease B subunit, produced as an enzymatically-inactive recombinant protein significantly decreases the level of H. felis infection in infected mice, in comparison to control infected mice.

The following examples are offered by way of illustration and are not intended to limit the invention in any manner.

A. Bacterial cultures and urease purification

The strain of Helicobacter pylori used in the study originates from a patient with a duodenal ulcer, and has been subcultured on BHI agarose plates to homogeneity. Helicobacter pylori is cultured in a suitable medium, typically, BHI (Brain-Heart Infusion) medium, containing 0.25% yeast extract and 10% fetal calf serum and supplemented with 0.4% Campylobacter selective complement (Skirrow supplement; Oxoid 69). The bacteria are incubated under microaerophilic conditions at 37°C in bottles that are sealed and shaken at 37°C for 2 to 3 days to produce a liquid culture. A culture may also be prepared in agarose plates consisting of BHI with 0.25% of yeast extract and 5% of sheep blood under microaerophilic conditions at 37°C for 3

days. The quantity of bacteria is determined by optical density of the BHI solution at 660 nm, with one optical density unit corresponding to 10^8 bacteria. Cultures on agarose plates are first resuspended in 154mM NaCl.

5 One currently preferred source of polyaminoacid displaying urease epitopes is purified urease, e.g., Helicobacter pylori urease obtained by following the general method of Dunn et al. J. Biol. Chem. 265, 9464-9469, modified as described below. Following culturing, the Helicobacter pylori is harvested in water, spun vortexed and spun again to produce a supernatant. The solution containing the urease activity of Helicobacter pylori (assessed by rapid urease test, see below) is then chromatographed on a CL-6B sizing column and the fractions which present a strong urease activity are pooled and dialysed overnight and again chromatographed on an anion exchanger gel. The fractions are eluted in increasing NaCl buffer and the collected fractions with a strong urease activity are individually submitted to a SDS gel followed by Coomassie staining. Two distinct bands corresponding to a molecular weight of about 63 and about
20 28 kDa are identified as urease. The fractions containing urease are pooled to give purified Helicobacter pylori urease having a purity in the region of 95% to 99%.

B. Oral immunization with urease purified from Helicobacter pylori

While it is preferred to employ purified Helicobacter pylori urease obtained as described above as the antigenic material, it will be understood that it is also possible to use, as the antigenic material, any urease or subunit of urease, either naturally occurring or obtained by recombinant DNA techniques, as well as digested fragment thereof, fusion proteins comprising the fragments or the whole urease, truncated urease constructs, or other peptide or protein preparations exhibiting urease epitopes which are capable of eliciting a protective immune response to Helicobacter infection (See below). Thus, it is possible to employ a urease having a substantial homology with respect to Helicobacter pylori urease and which is effective in raising a cross-protective immune response to Helicobacter. An example of such a urease is jack bean urease, which possesses about 70% homology with Helicobacter pylori urease. Although it was once thought that jack bean urease would be an effective antigen to prevent infection by Helicobacter, this is no longer believed to be the case. (See Chen, M. et al. "Failure of Immunization Against Helicobacter Using Jack Bean Urease," Acta Gastroenterol. Belg., Vol. 56, 94 (1993)). The invention is therefore not limited to the use of intact urease, and covers the

use of any polyaminoacid preparation which displays urease epitopes sufficiently homologous to Helicobacter pylori urease to generate a protective immunological response in a host to Helicobacter infection. A suitable urease must have sufficient
5 homology to H. pylori urease to elicit a protective immune response against Helicobacter infection. Typically, a urease having a homology of greater than 70%, for example, 80-90% homology, with respect to Helicobacter pylori urease, may be employed as the urease antigen in the invention.

10 A non-limiting list of sources of potentially useful urease preparations includes endogenous urease enzymes of the different Helicobacter species, urease from other bacteria such as Klebsiella pneumoniae or Proteus mirabilis, and, by analogy, any other urease which the condition that these ureases share cross-reactive epitopes with Helicobacter pylori urease. The urease genes of all the organisms mentioned above represent a potential tool for expressing recombinant urease products as a whole protein or as a part thereof.

20 A non-limiting list of potentially useful urease preparations includes peptides generated from purified urease (the sources are mentioned above), using physical and/or chemical cleavage procedures (i.e. CnBr) and/or proteolytic cleavage (using proteases e.g. V8-protease, trypsin or others); or

peptides synthesized chemically and retaining crossreactive epitopes with urease.

Other sources of potentially useful epitopes include epitopes identified by their crossreactivity with urease, as the
5 result of screening with anti-urease antibodies. These peptides can be naturally occurring peptides or peptides resulting from chemical synthesis. Furthermore such peptides can result from the expression of recombinant random oligonucleotides.

Another source of potentially useful epitopes includes epitopes similar to urease as a result of the generation of anti-
10 idiotypic antibodies to urease. Such anti-idiotypic antibodies, generated in any immunocompetent host, are obtained by immunization of this host with anti-urease antibodies, with the goal of generating antibodies directed against anti-urease
15 antibodies, which share structural homologies with urease.

The discussion herein focuses on the use of urease naturally produced by Helicobacter pylori (section B). However, it will be appreciated that the urease or subunits or constructs thereof mentioned above, capable of eliciting the desired
20 protective immune response, may be produced by recombinant DNA techniques well known in the art. The efficacy of particular preparations may be determined by routine administration using animal models, oral administration of the candidate vaccine, and

challenge with pathogen using a protocol substantially similar or identical to the procedure described below.

5 It will be recognized by those skilled in the art that other methods may be used to administer the Helicobacter urease peptides. For example, the urease peptides may be administered as part of a microsphere carrier formulation. Without limiting the type of microsphere carriers used, one example would be the administration of the urease peptides as part of a polylactide-coglycolide biodegradable microsphere carrier formulation.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
10185
10190
10195
10200
10205
10210
10215
10220
10225
10230
10235
10240
10245
10250
10255

virus-like particles, Rotavirus virus-like particles, or Ty particles, may be used to deliver the urease peptide. The aforementioned list is not meant to be limiting. Preferably, the live vector or the carrier system would be administered mucosally such that the recombinant urease peptides expressed by the live vector would be administered to a mucosal surface.

Tables 1 and 2 below and Figures 1-5 describe the results obtained when mice were orally immunized with purified Helicobacter pylori urease. In this first experiment, administration of the Helicobacter pylori antigen was carried out by orally administering to the mice Helicobacter pylori urease purified as described in section A, and coupled to hydroxyapatite crystals, used as a carrier to enhance M cell binding and uptake. Cholera toxin (Sigma) was given as a mucosal adjuvant. In this experiment, groups of female SPF BALB/c six-week old mice were each orally immunized with 30 μ g of purified Helicobacter pylori urease coupled to 1 mg of hydroxyapatite plus 10 μ g of cholera toxin adjuvant at day 0, 7, 14 and 21. The mice were then challenged twice with 10^8 H. felis, at day 28 and 30. For comparison purposes, similar female SPF BALB/c six-week old mice were orally immunized with whole Helicobacter pylori lysate (sonicate) and 10 μ g cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with H. felis. The Helicobacter pylori sonicate was prepared by collecting

Helicobacter pylori from cell cultures, pelleting by centrifugation and resuspending the pellet in 0.9% sodium chloride followed by sonication.

As a control, female SPF BALB/c six-week old mice were orally sham-immunized with 10 μ g of cholera toxin and 1 mg of hydroxyapatite at day 0, 7, 14 and 21. The mice were then challenged at day 28 and 30 with H. felis. All mice were housed and immunized in parallel. All mice subject to the study were sacrificed on day 35.

C. Oral immunization with recombinant urease subunits of Helicobacter pylori

Genes encoding the structural A and B subunits of Helicobacter pylori urease were obtained by polymerase chain reaction (PCR) cloning according to standard procedures, based on previously published sequences. (Clayton, C.L. et al. S. "Nucleotide Sequence of Two Genes from Helicobacter pylori Encoding for Urease Subunits" Nucleic Acid Res., vol. 18, 362 (1990)). These genes were inserted in a vector (named pEV40) designed for high expression and easy purification of foreign genes in E. coli. Briefly, the foreign gene is inserted downstream of a thermo-repressible promoter, and in frame of a sequence encoding for a repeat of six histidines. An ampR gene

is present on this vector for selection of transformants. Under the appropriate temperature conditions, the recombinant protein obtained is supplemented by six histidines at the N-terminal, which allow for a one-step affinity purification on a nickel column. Both Helicobacter pylori recombinant urease A and B subunits were expressed separately in E. coli, and purified on a nickel column to >95% purity.

While it is preferred to employ recombinant Helicobacter pylori urease obtained as described above as the antigenic material, it will be understood that it is also possible to use, as the antigenic material, any urease or subunit of urease obtained by recombinant techniques (e.g. fusion protein) expressing antigenic sites of urease, which is capable of eliciting a protective immune response to Helicobacter infection. Thus, it is possible to employ in a construct a urease gene having a substantial homology with respect to Helicobacter pylori urease and which is effective in raising a cross-protective immune response to Helicobacter. Examples of such a urease is jack bean urease, which possesses about 70% homology with Helicobacter pylori urease, or H. felis urease, which possesses about 88% homology with Helicobacter pylori urease. The invention is therefore not limited to the use of Helicobacter pylori urease genes and their gene products, and covers the use of any recombinant urease, or the subunits thereof, which is

0955739.094801
108460.6E25560

sufficiently close antigenically to generate a protective immunological response in a host to Helicobacter infection. The invention includes within its scope the use of any urease or subunit of urease, either naturally occurring or obtained by recombinant DNA techniques, as well as a digested fragment or peptide thereof, fusion protein comprising the whole urease, subunit or fragment thereof, or truncated urease construct which is effective in reducing the level of Helicobacter infection in an infected mammal. Typically, a recombinant urease having a homology of 70-95% homology, for example, 80-90% homology, with respect to Helicobacter pylori urease, may be employed as the recombinant urease antigen in the invention.

The discussion herein focuses on the use of recombinant Helicobacter pylori urease A and B subunits produced by E. coli (section C). However, it will be appreciated that recombinant urease or subunits or constructs thereof mentioned above, capable of eliciting the desired protective immune response, may be produced using other recombinant DNA techniques and other eukaryotic or prokaryotic expression vectors well known in the art.

Tables 3, 4 and 5 below and Figure 6 describe the results obtained when mice were orally immunized with recombinant Helicobacter pylori urease subunits produced in E. coli. In this experiment, administration of the Helicobacter pylori antigen was carried out by orally administering to the mice recombinant

108160-6625660
0955739-091801

Helicobacter pylori urease A or B subunits produced in E. coli and purified as described above, and coupled to hydroxyapatite crystals, used as a carrier to enhance M cell binding and uptake. Cholera toxin (Sigma) was given as a mucosal adjuvant. In this experiment, groups of female SPF BALB/c six-week old mice were each orally immunized with 30 μ g of recombinant Helicobacter pylori urease A subunit, coupled to 1 mg of hydroxyapatite plus 10 μ g of cholera toxin adjuvant at day 0, 8, 14 and 21. The mice were then challenged with 10^8 H. felis, at day 32, 34 and 36. For comparison purposes, similar female SPF BALB/c six-week old mice were orally immunized with 30 μ g of recombinant Helicobacter pylori urease B subunit coupled to hydroxyapatite plus 10 μ g cholera toxin at day 0, 8, 14 and 21. The mice were challenged three times, at day 32, 34 and 36, with H. felis. As a control, female SPF BALB/c six-week old mice were each orally sham-immunized with 10 μ g of cholera toxin and 1 mg of hydroxyapatite at day 0, 8, 14 and 21. The mice were then challenged at day 32, 34 and 36 with H. felis. All mice subject to the study were immunized and challenged in parallel. Animals were sacrificed on day 48 (12 days after challenge) or 10 weeks after challenge.

D. Analysis of gastric biopsies, blood, and intestinal secretions

Biopsies were taken from the stomach and blood was obtained from the heart. The intestines were removed and washed with 1mM PMSF (Boeringher) in PBS buffer to obtain intestinal secretions for ELISA analysis.

5 To evaluate protection against H. felis colonization, gastric biopsies from each animal were screened for the presence of H. felis by assessing rapid urease activity by the Jatrox HP test (Rohm Pharma), according to the supplier's directions. Briefly, gastric biopsies are immersed in 0.5 ml supplier's mixture of urea and phenol red, a pH indicator. A non-commercial version of the urea and phenol red mixture will be recognized by those skilled in the art and may contain, for example, Bacto yeast extract (0.1g), monopotassium phosphate (0.091g), disodium phosphate (0.095g), urea (20g), and Bacto phenol red (0.01g) at a final pH of 6.9 at 25°C. Urease activity generates ammonia and bicarbonate from urea, and is followed by the colorimetric change of the solution towards a higher absorbance at 550 nm. Urease activity was quantified by spectrophotometric analysis. Other methods of assaying urease activity will be recognized by those skilled in the art and may be found in, for example, Mobley, H.L.T. & Hausinger, R.P., "Microbial Ureases: Significance, Regulation, and Molecular Characterization," Microb. Reviews, Vol. 53, 85-108 (1989).

108760-525566
1055739-091901

20

Gastric biopsies of each animal included in the experiment described in section B were also cultured on BHI agarose plates, supplemented as above, for the detection of H. felis. After incubation for 3 to 10 days in microaerophilic conditions, the presence of H. felis was confirmed by Gram staining and determination of urease activity. As a very significant correlation was obtained for the detection of H. felis in gastric biopsies between urease tests and H. felis cultures during the first set of experiments (see Table 3), only gastric biopsy urease tests were performed for the detection of H. felis in the experiment described in section C. Detection of H. felis was confirmed by microscopy by two independent investigators, using two different colorations (acridine orange and cresyl violet).

Blood samples were allowed to clot for 3 hours at RT (room temperature), and serum harvested and frozen at -20°C, until further analysis. Intestinal secretions were spun for 5 minutes at 4°C to remove debris, and kept frozen at -20°C. Serum and intestinal samples of each animal were analyzed by ELISA for evaluation of anti-Helicobacter activity, according to standard procedures. Briefly, 96-well plates were coated with a sonicate of Helicobacter pylori, followed by saturation with 5% fat-free milk. Samples were serially diluted from 1:1 to 1:1000 and incubated overnight at 4°C on ELISA plates. Biotinylated anti-

105T60-6E25660

mouse IgG (serum) and anti-mouse IgA, followed by streptavidin-Horseradish peroxidase was used for the determination of the antibody levels.

The results of H. felis challenges following immunizations with purified Helicobacter pylori urease are set out in Tables 1-3 and Figures 1-4 and the results of H. felis challenges following immunizations with recombinant Helicobacter pylori urease A and B subunits are set out in Tables 4-6 and Figures 5 and 6.

09055739.091801
TOBT60" 6E/55660

TABLE 1

mouse number	Immunization	urease test	culture Gram	Immunoglobulins			
				Serum		Intestinal secretion	
		12h		Ig	Ig G	Ig	IgA
1	Urease+HF	+	<i>H felis</i>	27	0	25	258
2	Urease+HF	0	0	264	273	221	246
3	Urease+HF	0	0	84	44	318	354
4	Urease+HF	+	<i>H felis</i>	81	42	12	5
5	Urease+HF	0	0	98	137	126	234
6	Urease+HF	+	0	968	2093	31	22
7	Urease+HF	0	0	98	0	96	34
8	Urease+HF	0	0	247	1010	214	128
9	Urease+HF	0	0	N.D.	N.D.	48	23
10	Urease+HF	0	0	50	0	124	99
11	Urease	0	0	319	205	44	53
12	Urease	0	0	14	0	86	87

13	Urease	0	0	0	0	0	0
14	Urease	0	0	0	0	43	61
15	Urease	0	0	58	0	110	127
16	Urease	0	0	140	63	21	37
17	Urease	0	0	84	240	114	280
18	Urease	0	0	N.D.	N.D.	93	148
19	Urease	0	0	45	0	135	216
20	Urease	0	0	261	197	161	261
21	CT+HF	0	0	0	0	0	2
22	CT+HF	+	<i>H felis</i>	63	0	310	303
23	CT+HF	+	<i>H felis</i>	90	0	N.D.	N.D.
24	CT+HF	+	<i>H felis</i>	31	0	150	192
25	CT+HF	+	<i>H felis</i>	197	250	250	440
26	CT+HF	+	<i>H felis</i>	105	135	214	138
27	CT+HF	+	<i>H felis</i>	140	47	109	55
28	CT+HF	+	0	0	0	16	15
29	CT+HF	+	<i>H felis</i>	0	0	0	0

30	CT+HF	+	<i>H felis</i>	N.D.	N.D.	N.D.	N.D.
31	HP sonicate+HF	+	<i>H felis</i>	0	0	76	103
32	HP sonicate+HF	+	<i>H felis</i>	77	0	11	33
33	HP sonicate+HF	+	<i>H felis</i>	549	748	57	36
34	HP sonicate+HF	0	0	660	153	180	286
35	HP sonicate+HF	+	<i>H felis</i>	730	192	0	5
36	HP sonicate+HF	+	<i>H felis</i>	32	0	5	64
37	HP sonicate+HF	0	0	400	400	312	1149
38	HP sonicate+HF	+	<i>H felis</i>	1007	1360	149	26
39	HP sonicate+HF	0	0	220	186	133	122
40	HP sonicate	0	0	873	1016	352	514
41	HP sonicate	0	0	727	899	126	191
42	HP sonicate	0	0	109	68	44	83
43	HP sonicate	0	0	147	949	167	97
44	HP sonicate	0	0	845	1094	246	64
45	HP sonicate	0	0	1217	1198	210	157
46	HP sonicate	0	0	81	0	256	218

47	HP sonicate	0	0	329	210	241	276
48	HP sonicate	0	0	1049	737	197	211

In Table 1, which refers to the experiment described in section B, "h" means hour, "Ig" means immunoglobulin, "ND" means "not determined", "urease + HF" means that the mice were immunized with urease (coupled to hydroxyapatite, with cholera toxin) and then challenged with H. felis, "urease" means that the mice were immunized with urease (coupled to hydroxyapatite, with cholera toxin) and not challenged, "CT+HF" means that the mice were sham-immunized with cholera toxin and challenged with H. felis, "HP sonicate + HF" means that the mice were immunized with Helicobacter pylori sonicate with cholera toxin and challenged by H. felis, and "HP sonicate" means that the mice were immunized with Helicobacter pylori sonicate with cholera toxin and not challenged. In Table 1, the numbers for the antibody results are given as a measure of absorbance at 595 nm multiplied by 1000. The background measured in absence of the antibodies, was subtracted.

The results of experiment described in section B obtained on the basis of the gastric biopsies urease tests and on Gram staining of H. felis cultures are set out in Table 2. Infec-

tion was defined by mice with one or more markers of colonization by H. felis, including urease test or Gram staining of cultures.

TABLE 2

<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>	
5 Urease	<u>H. felis</u>	3/10 (30%)	7/10 (70%)	*
Sonicate	<u>H. felis</u>	6/9 (66%)	3/9 (33%)	**
CT	<u>H. felis</u>	9/10 (90%)	1/10 (10%)	

* p=0.0198 (two tailed Fisher exact test) compared to CT control

** p=0.303 (two tailed Fisher exact test) compared to CT control

It will be seen from the results set out in Tables 1 and 2 that statistically significant protection against H. felis challenge is obtained with oral immunization using Helicobacter pylori urease as compared to that obtained using either Helicobacter pylori sonicate or cholera toxin. Referring to Table 2, it will be seen that from a total of 10 immunized animals, only 3 became infected, as compared to 6 of the animals immunized with Helicobacter pylori sonicate and 9 of the animals immunized with cholera toxin. Table 2 shows that 70% of the animals were protected from challenge by H. felis as compared to 33% of the

animals immunized with Helicobacter pylori sonicate and 10% of the animals immunized with cholera toxin and then subjected to H. felis challenge. In other words, 90% of the control mice exposed to H. felis became infected by that pathogen whereas, in contrast, in mice immunized with Helicobacter pylori urease 28 days before exposure to H. felis, the infection rate was only 30%. This represents a significant reduction in infection ($p=0.0198$ in the Fisher exact test, as compared to the control mice). When the mice were orally immunized with Helicobacter pylori sonicate, the infection rate was 67% (not significant versus the control). The protection obtained using Helicobacter pylori urease is unexpected and could not have been predicted on the basis of the results observed using Helicobacter pylori sonicate.

Referring to Figures 1-4, Figure 1 represents graphically the results of tests for antibodies in serum (IgG) and intestinal secretion (IgA) in mice not protected after immunization with urease. These are mice numbers 1, 4 and 6 appearing in Table 1, and constitute Group A. Figure 2 shows the antibody responses of mice that were protected after immunization with urease (Group B), i.e. mice 2, 3, 5 and 7-10.

Figures 3 and 4 relate to the results obtained with mice numbers 31-39. Figure 3 (Group C) depicts antibody responses of mice not protected after immunization with Helicobacter pylori sonicate (mice numbers 31, 32, 33, 35, 36 and 38) and Figure 4

(Group D) depicts the antibody responses of mice protected after immunization with Helicobacter pylori sonicate (mice numbers 34, 37 and 39). It is of interest to note with respect to Figures 3 and 4 that the IgA antibody responses (but not IgG) are higher in the mice exhibiting protection than in the mice that are not protected, suggesting a correlation between protection and IgA response. Serum IgG responses did not exhibit a correlation. Mucosal IgA but not serum IgG are known to play a role in protection against bacterial infections of the gut. (McGhee, J.R and Kyono, H. "New Perspectives in Vaccine Development: Mucosal Immunity to Infections" Infect Agents Dis., vol. 2, 55-73 (1993)).

The results of the correlation between the detection of H. felis in gastric biopsies by urease tests and by cultures are set out in Table 3.

TABLE 3

	Urease Test +	Urease Test -	Total
<u>H. felis</u> culture (+)	16	0	16
<u>H. felis</u> culture (-)	2	30	32
Total	18	30	48

Two-tailed Fisher's Exact Test: $p < 0.00001$

Table 3 shows that a very significant correlation exists between the results of urease tests performed on gastric biopsies and the identification of H. felis by cultures. All animals shown in Table 3 that were positive for urease activity, were H. felis positive by histopathology. As cultures detected less often H. felis infection than urease tests, urease tests were preferred for the diagnosis of H. felis infection in mice in the next experiments, due to its better sensitivity. This approach allowed the duplication of urease tests with larger fragments of the stomach of each mouse, and a further increase in the sensitivity of the urease test. Furthermore, the use of the method with the highest sensitivity prevent an overestimation of the protection obtained by the vaccine preparation to be tested. When positive culture is used as the standard for infection, the protection induced after urease immunization during the experiment depicted in section B is as significant as with the combined use of urease test and culture ($p=0.021$ versus $p=0.019$).

The results of the experiments described in section C (recombinant urease subunits), obtained on the basis of the gastric biopsies urease tests, are set out in Table 4, 5 and 6 and depicted in Figure 6.

TABLE 4

immunization	mice n°	Urease test	Infection
CT Sacrificed 12 days post challenge	20	0.49	+
	21	0.31	+
	22	0.62	+
	23	0.67	+
	24	0.55	+
	50	0.50	+
	51	0.37	+
	52	0.29	+
	53	0.79	+
	54	0.32	+
ure A + HAP + CT Sacrificed 12 days post challenge	40	0.67	+
	41	0.48	+
	42	0.42	+
	43	0.65	+
	44	0.56	+
	45	0.52	+
	46	0.33	+
	47	0.63	+
	48	0.22	+
	49	0.37	+
ure B + HAP + CT Sacrificed 12 days post challenge	25	0.15	-
	26	0.07	-
	27	0.03	-
	28	0.64	+
	29	0.13	-
	30	0.02	-
	31	0.66	+

	32	0.00	-
	33	0.79	+
	34	0.15	-
ure A + HAP + CT	68	0.00	-
Sacrificed 10 weeks	69	0.07	-
post challenge	70	0.42	+
	71	0.00	-
	72	0.00	-
ure B + HAP + CT	73	0.37	+
Sacrificed 10 weeks	74	0.00	-
post challenge	75	0.37	+
	76	0.00	-
	77	0.00	-
	78	0.00	-
	79	0.39	+
	80	0.00	-
	81	0.37	+
	82	0.00	-

In Table 4, "CT" means cholera toxin; "UreA" means recombinant Helicobacter pylori urease A subunit; "Ure B" means recombinant Helicobacter pylori urease B subunit; and "HAP" means hydroxyapatite crystals. Mice 20 to 54 were sacrificed 12 days post challenge and mice 68 to 82 10 weeks (106 days) post challenge. The results of the urease test performed from biopsies of the stomach of each animal are expressed as OD values at 550 nm. The O.D. value for the assay solution alone (0.075

0055739.091801

O.D.) was subtracted as background from the O.D. value obtained for each mouse. The positive and negative signs depicts the final status of infection of each animal, according to the positivity or negativity of the urease test for detection of H. felis. Positivity: OD₅₅₀ values >0.2. The 0.2 value was chosen to define a positive infection because at this value, the color change to the Jatrox solution could be observed with the naked eye.

TABLE 5: Protection as measured 12 days post challenge

<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>
Urease A subunit	<u>H. felis</u>	10/10 (%)	0/10 (0%)
Urease B subunit	<u>H. felis</u>	3/10 (30%)	7/10 (70%) *
CT	<u>H. felis</u>	10/10 (100%)	0/10 (0%)

* p=0.0031 (two tailed Fisher exact test) compared to CT control

TABLE 6: Protection as measured 10 weeks post challenge

<u>Immunization</u>	<u>Challenge</u>	<u>% infected</u>	<u>% protected</u>
Urease A subunit	<u>H. felis</u>	1/5 (20%)	4/5 (80%)*
Urease B subunit	<u>H. felis</u>	4/10 (40%)	6/10 (60%)**

* p=0.003 (two tailed Fisher exact test) compared to CT control

** p=0.01 (two tailed Fisher exact test) compared to CT control

It will be seen from the results set out in Tables 4, 5 and 6 that statistically significant protection against H. felis challenge is obtained with oral immunization using recombinant Helicobacter pylori urease B subunit as compared to that obtained using either recombinant Helicobacter pylori urease A subunit or cholera toxin. Referring to Table 4, it will be seen that, 12 days post challenge, from a total of 10 immunized animals, only 3 were found infected in the urease B subunit group, as compared to all 10 animals immunized with Helicobacter pylori A subunit of urease and 10 out of 10 of the animals immunized with cholera toxin. Table 4 shows that 70% of the animals were protected from challenge by H. felis as compared to 0% of the animals immunized with Helicobacter pylori urease A subunit and 0% of the animals immunized with cholera toxin and then subjected to H. felis challenge. In other words, 100% of the control mice challenged

with H. felis became infected whereas, in contrast, in mice immunized with recombinant Helicobacter pylori urease B subunit the infection rate was only 30%. This represents a significant reduction in infection ($p=0.0031$, Fisher exact test) as compared to the control mice. The fact that the protection observed with Helicobacter pylori urease is entirely conferred by immunization with the B subunit of urease, and that the A subunit has no such effect, was not expected on the basis of our experiment with purified urease. This definition of the rôles of the 2 structural subunits of urease in the development of a protective immune response is therefore novel. The protection obtained using recombinant urease subunits, which are enzymatically inactive, also teaches that non toxic forms of urease can be used as oral vaccine against Helicobacter infection. Furthermore these results strongly suggest that recognition of the active site is not required for protection, as an inactive urease B subunit is very unlikely to induce antibodies that will recognize and inhibit the catalytic site of native urease.

Referring to Table 6, it will be seen that, when mice are sacrificed 10 weeks post infection, 60% (6 mice out of 10) of the animals immunized with urease B subunit and 80% (4 mice out of 5) of the animals immunized with Helicobacter pylori urease A subunit were protected against H. felis infection. The fact that protection obtained through immunization with urease B subunit

lasts over time and that immunization with urease A induces a protection which is delayed compared to the one induced by urease B subunit could not be expected from our experiment with purified urease or with other experiment performed earlier. The fact that urease A subunit immunization confers protection definitively proves that recognition of the active site is not required for protection.

Figure 6 summarizes results obtained after oral immunization with recombinant urease A and B' subunits (described in Table 5 and 6).

A second set of mice was immunized and assayed for infection with Helicobacter felis at 10 weeks post challenge according to the protocol described in Section C. In this example, twelve mice were sham immunized with cholera toxin alone, twelve mice were immunized with recombinant ure A subunit, and ten mice were immunized with recombinant ure B subunit. The mean urease activity level found in stomach samples mice that were immunized with ure B, but not infected (uninfected mice controls), was 0.045, and this amount was subtracted as background from each O.D. value obtained. Mice were considered to be infected when the O.D. value was greater than twice the standard deviation of values obtained with the uninfected mice controls; the standard deviation was 0.022. The data obtained from this experiment is presented in Table 7.

10955739.091601
10955739.091601

TABLE 7

immunization	mice no.	Urease test	Infection
CT Sacrificed 10 weeks post challenge	135	0.40	+
	136	0.28	+
	137	0.25	+
	138	0.10	+
	139	0.10	+
	140	0.34	+
	141	0.41	+
	142	0.36.	+
	143	0.46	+
	144	0.40	+
	145	0.40	+
	146	0.51	+
ure A + HAP + CT Sacrificed 10 weeks post challenge	161	0.12	+
	162	0.47	+
	163	0.00	-
	164	0.00	-
	165	0.02	-
	166	0.01	-
	167	0.01	-
	168	0.37	+
	169	0.00	-
	170	0.39	+
	171	0.47	+
	172	0.00	-
ure B + HAP + CT Sacrificed 10 weeks	151	0.00	-
	152	0.00	-

post challenge	153	0.00	-
	154	0.03	-
	155	0.00	-
	156	0.00	-
	157	0.02	-
	158	0.00	-
	159	0.01	-
	160	0.00	-

Using this alternative method of analysis, the data presented in Table 4 was reanalyzed along with the data in Table 7. Instead of subtracting the background O.D. value obtained with the urease assay solution alone (0.075 O.D.), the mean urease level obtained from the uninfected mice controls was used as the background level.

The mean urease level obtained from uninfected mice sacrificed at twelve days was 0.089. Mice numbered 20-54 were considered infected when the O.D. value was greater than twice the standard deviation of values obtained with the uninfected mice controls; the standard deviation was 0.008. The background level subtracted from the O.D. values of mice sacrificed at ten weeks was 0.045, and mice were considered infected when the O.D. value was greater than 0.044.

Using this alternative method of analysis, at twelve days post challenge, no effect was seen after sham immunization

or after immunization with the ure A subunit. However, only a low-grade infection (O.D. value < 0.22) was observed in 70% of the animals immunized with the ure B subunit ($p < 0.02$, Mann-Whitney U-test, compared to sham immunized control). When mice were sacrificed ten weeks post challenge, 59% (10/17) of the mice immunized with the ure A subunit were protected against H. felis infection ($p = 0.0019$, two-tailed Fisher's exact test, when compared to sham-immunized mice). Furthermore, 80% (16/20) of the mice immunized with the ure B subunit were protected against H. felis infection ($p = 0.00002$, two-tailed Fisher's exact test, when compared to control, sham-immunized mice). Under this alternative analysis, the ure A subunit also elicits protective immunity. Under either method of analysis, the results obtained demonstrate that immunization with recombinant Helicobacter pylori urease subunits elicits protective immunity against Helicobacter infection.

EXAMPLES

The invention will now be further described by reference to the following non-limiting examples.

a) The Bacterial Strains

H. felis was provided by J. Fox (Division of Comparative Medicine, Mass. Institute of Technology, Boston,

USA). Helicobacter pylori was isolated from patients with ulcer disease (CHUV, Lausanne, Switzerland).

b) Bacterial Cultures

5 Liquid Culture - Bacteria were cultured on BHI (Brain-Heart Infusion, BioMerieux) liquid medium containing 0.25% of yeast extract (Difco) and 10% of fetal calf serum (Inotech) supplemented with 0.4% of Campylobacter selective complement (Oxoid). The bacteria were incubated under microaerophilic conditions at 37°C and shaken at 37°C for 2 to 3 days.

108160-6475660
15 Frozen Culture - The bacteria were cultured in liquid media, then assayed for urease activity, and evaluated for morphology by Gram staining and for motility by microscopy. The bacteria were then centrifuged and resuspended at a concentration of 30 O.D. per ml in BHI plus 20% glycerol and frozen at -80°C. Just before use, frozen stocks were thawed on ice, washed in 20 ml BHI, centrifuged, and resuspended at a concentration of 1 O.D. to 1.5 O.D. per 200 µl in 5mM NaHCO₃.

20 Culture on Agarose plates - The bacteria were cultured on agar plate consisting on BHI with 0.25% of yeast extract and 5% of sheep blood under microaerophilic conditions at 37°C for 3 days.

Quantification - The quantity of bacteria was determined by the optical density of the BHI solution at 660 nm (1 optical density unit (O.D.) corresponding to 10^8 bacteria). The number of viable bacteria is measured on the number of colony forming units.

c) Preparation of sonicates

Helicobacter pylori was collected from 31 blood agar plates in 0.15 M NaCl and spun 5 minutes at 1400g at 4°C. The pellet was resuspended in 3 ml of NaCl and sonicated for 4 minutes. The amount of proteins was evaluated by a Bradford assay (BioRad Kit according to supplier).

d) Coupling of immunogen to Hydroxyapatite

Immunogen (urease or subunit thereof) was incubated for 1 hour at 4°C with hydroxyapatite. 1.0 mg of hydroxyapatite was used for 30 µg of immunogen per mouse. At the end of the incubation, 10 µg of cholera toxin was added in a final volume of 200 µl PBS.

e) Challenge with Helicobacter felis

Mice were lightly anesthetized prior to intragastric challenge with Helicobacter felis. H. felis in a total volume of

200 μ l were delivered to the stomachs of the respective mice using silicon tubing attached to a hypodermic syringe.

EXAMPLE 1

a) Extraction

5 Helicobacter pylori from 30 blood agar plates was harvested in 0.15 M NaCl on ice. The solution was spun 5 minutes at 1400 g at 4°C. The pellet was resuspended in 20 ml of H₂O and vortexed for 45 seconds (maximum speed). The extract was then spun 20 minutes at 6700 g at 4°C. The supernatant was recovered and the quantity of protein was evaluated (see "Quantification" above) and precipitated with 70% of ammonium sulfate.

b) Purification of urease

15 The solution was chromatographed on a Sepharose CL-6B column (Pharmacia) with PBS (phosphate buffered saline) as mobile phase. The 22 collected fractions which presented a strong urease activity were pooled and dialyzed overnight at 4°C against 3 liters of PEB (20 mM phosphate buffer, pH 7) and then chromatographed on a Q Sepharose fast flow (Pharmacia) with PEB as mobile phase. The fractions were eluted by 0 to 500 mM NaCl gradient.

20 Ten of the collected fractions with a strong urease activity were individually subjected to an SDS gel followed by a Coomassie staining. The 6 fractions presenting 2 distinct bands

corresponding to MW-63 and -28 KDa were pooled and were considered as the purified urease.

EXAMPLE 2 (see also section B)

5 Mice employed in the immunization studies were lightly anesthetized with ether prior to intragastric immunization. And then, sonicate preparation or purified urease, hydroxyapatite and cholera toxin was suspended in PBS and 200 μ l were delivered to the stomach of the respective mice using a polyethylene tubing attached to a hypodermic syringe. This procedure will be referred to as oral immunization.

Three oral immunization protocols were evaluated. These are described below.

Protocol B1 - Vaccination with purified urease

15 Female BALB/c six-week old mice (20) were orally immunized with 30 μ g of purified of Helicobacter pylori urease and 1 mg of hydroxyapatite and 10 μ g of cholera toxin at day 0, 7, 14 and 21. Ten mice were challenged at day 28 and 30 with 5×10^7 and 10^8 H. felis from liquid culture.

Protocol B2 - Vaccination with Helicobacter sonicates

Female BALB/c six-week old mice (20) were orally immunized with 2 mg of Helicobacter pylori sonicate solution at day 0, 7, 14 and 21. Ten mice were challenged at day 28 and 30 with 5×10^7 and 10^8 H. felis.

Protocol B3 - Control

Female BALB/c six-week old mice were orally immunized with 1 mg hydroxyapatite and 10 μ g of cholera toxin at day 0, 7, 14 and 21. The mice were challenged at day 28 and 30 with 5×10^7 and 10^8 H. felis.

At day 35 all mice were sacrificed and biopsies from the stomach were taken as well as intestinal secretions and blood.

Protection and evaluation

To evaluate protection, biopsies were screened for the urease activity by the Jatrox HP test (Rohm Pharma) according to the instructions of the supplier. The urease is quantified by a spectrophotometric measurement at 550nm. The biopsies were also cultured for the presence of H. felis and the presence of H. felis was determined by Gram staining. Gastric antral biopsies were homogenized and diluted (1:10 and 1:1000) in 0.15 M NaCl and

plated onto blood agar plates and incubated under microaerophilic conditions at 37°C for 4 to 10 days.

ELISA

Intestinal secretions and blood were analyzed by ELISA for the evaluation of antibody titer. The ELISA was carried out as follows. Polystyrene plates (96 wells) were coated with 1 μ g/well of purified urease at 37°C for 2 hrs. Non specific binding sites were blocked with 5% powdered milk in PBS 0.1% Tween at 37°C for 30 minutes. The plates were washed once with PBS 0.1% Tween. Blood samples were tested at dilution 1:1000 and intestinal secretions 1:1. 100 μ l of each sample were added to the antigen coated plates. After 2 hrs of incubation, plates were washed 3 times with PBS containing 0.1% Tween. Anti-mouse biotinylated whole antibody from goat and anti-mouse IgA, IgG and IgM biotinylated (Amersham) were added (100 μ l) at dilution 1:500 except for IgA (1:250) and incubated at 37°C for 1 hr. The plates were washed 3 times with PBS 0.1% Tween and 100 μ l of 1:1000 dilution of streptavidin Horseradish peroxidase in PBS containing 0.1% Tween were added and incubated at 37°C for 30 minutes. The plates were washed 3 times and 50 μ l of 1:50 dilution of o-phenyl-diamine in citrate buffer pH 5.0 with 1 μ l/ml of 30% H₂O₂ were added and incubated at room temperature

for 20 minutes. The absorbance at 495 nm was measured in each well.

EXAMPLE 3 (see also section C)

Mice employed in the immunization studies were lightly anesthetized with ether prior to intragastric immunization. Then, 30 μ g recombinant Helicobacter pylori urease A or B subunit produced in E. coli, bound with hydroxyapatite, and supplemented with cholera toxin was suspended in PBS and 200 μ l were delivered to the stomach of the respective mice using a polyethylene tubing attached to a hypodermic syringe. This procedure will be referred to as oral immunization.

Three oral immunization protocols were evaluated. These are described below.

Protocol C1 - Vaccination with recombinant urease A subunit

Female BALB/c six-week old mice (10) were orally immunized with 30 μ g of purified recombinant Helicobacter pylori urease A subunit and 1 mg of hydroxyapatite and 10 μ g of cholera toxin at day 0, 8, 14 and 21. Ten mice were challenged at day 32, 34 and 36 with 10^8 H. felis from liquid culture.

Protocol C2 - Vaccination with recombinant ur as B subunit

Female BALB/c six-week old mice (10) were orally immunized with 30 µg of recombinant Helicobacter pylori urease B subunit and 1 mg of hydroxyapatite and 10 µg of cholera toxin at day 0, 8, 14 and 21. Ten mice were challenged at day 32, 34 and 36 with 10⁸ H. felis from liquid culture.

Protocol C3 - Control

Female BALB/c six-week old mice were orally immunized with 1 mg of hydroxyapatite and 10 µg of cholera toxin at day 0, 8, 14 and 21. The mice were challenged at day 32, 34 and 36 with 10⁸ H. felis.

At day 42, or at day 106, mice were sacrificed and multiple biopsies from the stomach were taken.

Protection and evaluation

To evaluate protection, biopsies of the corpus and antrum of the stomach were screened for urease activity by the Jatrox HP test (Rohm Pharma) according to the instructions of the supplier. The urease is quantified by a spectrophotometric measurement at 550nm. The total of corpus and antrum OD values were added to obtain a final OD value for each mouse.

EXAMPLE 4

To determine whether immunization with urease peptides would be an effective treatment of Helicobacter infection animals, mice were first challenged with H. felis and then immunized with Helicobacter pylori ure B subunit. The utility of immunization with Helicobacter pylori ure B subunit to treat Helicobacter infection is demonstrated in both Example 4 and Example 5.

a) Infection of mice with H. felis

Female BALB/c six-to-eight week old mice were challenged with H. felis at days 1 and 3 with 1 O.D. of frozen culture. The mice were challenged at day 5 with 1.53 O.D. of liquid H. felis culture.

b) Vaccination with recombinant Helicobacter pylori urease B subunit

Eight H. felis infected mice were orally immunized with 30 μ g of recombinant Helicobacter pylori urease B subunit, 1 mg of hydroxyapatite and 10 μ g of cholera toxin (holoenzyme obtained from Calbiochem) at days 23, 30, 37, and 44.

c) Controls

Ten H. felis infected mice were orally immunized with 1 mg of hydroxyapatite and 10 μ g of cholera toxin at days 23, 30, 37, and 44. These mice were designated as "sham" immunized.

Seven H. felis infected mice were not subject to immunization.

d) Sacrifice and Evaluation

At day 70 all mice were sacrificed. The stomachs were removed and cut in half longitudinally. To evaluate protection, one-half of the stomach of each mouse (both corpus and antrum sections) was screened for urease activity 3-4 hours post-sacrifice, by the Jatrox HP test (Rohm Pharma) according to the instructions of the supplier. The urease was quantitated by a spectrophotometric measurement at 550 nm. Ten naive (uninfected) BALB/c mice served as controls. These mice were sacrificed at day 70 and the stomach samples evaluated for urease activity to determine the background level of urease activity. The mean background value, 0.033 O.D., was subtracted from the O.D. value for each mouse. The standard deviation in urease activity among the naive mice was 0.025 O.D. The results of the urease assays are set out in Table 8 and depicted in Figure 7. In Figure 7, the dashed line indicates the O.D. value for twice the standard deviation in urease activity among the naive mice.

TABLE 8

Treatment	Mouse No.	Urease test	Infection
Untreated	1	0.42	+
	2	0.36	+
	3	0.41	+
	4	0.33	+
	5	0.38	+
	6	0.52	+
	7	0.48	+
Sham (HAP+CT only)	8	0.51	+
	9	0.48	+
	10	0.61	+
	11	0.44	+
	12	0.51	+
	13	0.48	+
	14	0.44	+
	15	0.58	+
	16	0.48	+
	17	0.53	+
Ure B + HAP + CT	18	0.36	+
	19	0.01	-
	20	0.43	+
	21	0.07	+/-
	22	0.39	+
	23	0.10	+/-
	24	0.38	+
	25	0.06	+/-

In Table 8, "ure B" means recombinant Helicobacter pylori urease B subunit as referenced above; "CT" means cholera toxin; and "HAP" means hydroxyapatite crystals; "untreated" means the mice were challenged with H. felis, but received no subsequent immunization. The results of the urease test performed

from biopsies of the stomach of each animal are expressed as O.D. values at 550 nm. The background value subtracted from the O.D. value for each mouse was 0.033. The infection is rated as positive (indicated by a "+") where the O.D. value at 550 nm is over twice the standard deviation found with the naive mouse controls (0.05 O.D.). A "+/-" indicates that the O.D. value is greater than background but less than 0.08, and that the value is reduced compared to the sham immunized controls.

It will be seen from the results set out in Table 7 and in Figure 7 that administration of the Helicobacter pylori ure B subunit to mice infected with H. felis resulted in the clearance of infection in one out of eight mice. According to another, less conservative, interpretation of these results, defining a positive result as a value greater than 0.2, four out of eight mice (numbers 2, 4, 6, and 8) treated with ureB would be found to be protected from infection. However, under either of the two definitions of a positive result these mice exhibited a reduced urease activity as compared to the untreated and the sham-immunized mice. Using the 0.05 value to define infection, the level of infection in ure B immunized mice as compared to sham immunized mice exhibits a statistically significant (p value of less than 0.004) reduction in the level of infection. Thus the results of Example 4 indicate that administration of ure B to

mice infected with H. felis results in a reduced level of infection.

EXAMPLE 5

a) Infection of mice with H. felis

5 Female BALB/c five-to-eight week old mice were challenged with H. felis at day 1 with 1 O.D. of liquid culture. The mice were challenged at day 3 with 0.8 O.D. H. felis cultured on agarose plates. The mice were challenged at day 5 with 1 O.D. of H. felis cultured on agarose plates.

09955739.091801
108160.68755660

b) Vaccination with recombinant *Helicobacter pylori* urease B subunit

H. felis infected mice were orally immunized with 30 μ g of recombinant *Helicobacter pylori* urease B subunit, 1 mg of hydroxyapatite and 10 μ g of cholera toxin (holoenzyme obtained from Calbiochem) at days 23, 30, 37, and 44. These mice were designated as "immunized."

c) Controls

H. felis infected mice were orally immunized with 1 mg of hydroxyapatite and 10 μ g of cholera toxin at days 23, 30, 37, and 44. These mice were designated as "sham" immunized.

d) Sacrifice and Evaluation

At day 62, ten immunized mice and six sham immunized mice were sacrificed. At eight weeks after the last immunization, ten immunized and eight sham immunized mice were sacrificed. Stomach samples were screened for urease activity as described in Example 4. The mean background found in 29 naive BALB/c mice, sacrificed at various times, was 0.04 O.D. This level of activity was used as a background measurement and subtracted from the O.D. value for each mouse. The standard deviation in urease activity among the naive mice was 0.02 O.D. The results of the urease assays are set out in Table 9 and

depicted in Figure 8. In Figure 8, the dashed line indicates the O.D. value for twice the standard deviation in urease activity among naive mice.

e) Analysis of Blood and Fecal Samples

Blood Sample Collection

Blood samples were allowed to clot for 3 hours at room temperature, and serum harvested and frozen at -20°C, until further analysis.

Fecal Pellet Collection

Fresh fecal pellets (3-8) were collected in 1.5 ml eppendorf tubes containing 500 μ l of PBS +5% non-fat dry milk + protease inhibitors (AEBSF 0.2 mM, Aprotinin 1 μ g/ml, Leupeptin 10 μ M, Bestatin 3.25 μ M). The samples were frozen at -20°C until use. They were then thawed on ice, mashed, and centrifuged at 10,000g for 10 minutes at 4°C to yield a supernatant free of debris, and yellowish brown in appearance.

ELISA

Serum and fecal pellet samples of each animal were analyzed by ELISA for evaluation of anti-urease activity, according to standard procedures.

Fecal pellets and blood were analyzed by ELISA for the evaluation of antibody titer. Polystyrene plates (96 wells) were coated with 0.5 μ g/well of purified recombinant urease at 37°C for 2 hrs. Non specific binding sites were blocked with 5% powdered milk in PBS containing 0.1% Tween at 37°C for 30 minutes. The plates were washed once with PBS containing 0.1% Tween. Blood samples were tested at a dilution of 1:200 and fecal pellets at a dilution of 1:1. 100 μ l of each sample were added to the antigen coated plates. After 2 hrs of incubation, plates were washed 3 times with PBS containing 0.1% Tween. Anti-mouse biotinylated whole antibody from goat (Amersham) and anti-mouse IgA coupled to Horseradish peroxidase (Serotec) were added (100 μ l) at a dilution of 1:500 and incubated at 37°C for 1 hr. The plates were washed 3 times with PBS containing 0.1% Tween and, for blood samples only, 100 μ l of a 1:500 dilution of streptavidin and Horseradish peroxidase in PBS containing 0.1% Tween were added and incubated at 37°C for 30 minutes. The plates were washed 3 times and 50 μ l of a 1:50 dilution of o-phenyl-diamine in citrate buffer, pH 5.0, with 1 μ l/ml of 30% H₂O₂ were added and incubated at room temperature for 20 minutes. The absorbance at 495 nm was measured in each well.

For the evaluation of total IgAs in fecal pellets, the ELISA was carried out as described above except that the plates

were coated with 1 μ g/ml of goat anti-mouse IgA (SIGMA) and the fecal supernatant was tested at a 1:200 dilution.

TABLE 9

Sacrifice	Treatment	Mouse No.	Urease test	Infection
2.5 weeks	Sham (HAP+CT only)	1	0.58	+
		2	0.51	+
		3	0.50	+
		4	0.54	+
		5	0.47	+
		6	0.36	+
	Ure B + HAP + CT	7	0.02	-
		8	0.06	+
		9	0.01	-
		10	0.00	-
		11	0.02	-
		12	0.02	-
		13	0.02	-
		14	0.44	+
		15	0.00	-
		16	0.03	-
8 weeks	Sham (HAP+CT only)	17	0.30	+
		18	0.27	+
		19	0.29	+
		20	0.20	+
		21	0.27	+
		22	0.27	+
		23	0.37	+
		24	0.31	+
	ure B +HAP+CT	25	0.26	+
		26	0.30	+
		27	0.00	-
		28	0.17	+
		29	N.D.	
		30	0.00	-

Sacrifice	Treatment	Mouse No.	Urease test	Infection
		31	0.00	-
		32	0.00	-
		33	0.00	-
		34	0.00	-

TABLE 10

Sacrifice	Treatment	% cleared
2.5 weeks	ure B	8/10 (80%)*
	sham	0/6
8 weeks	ure B	6/9 (67%)**
	sham	0/9

* p = 0.007 (two-tailed Fisher exact test) compared to sham control

** p = 0.014 (two-tailed Fisher exact test) compared to sham control

In Tables 9 and 10, "ure B" means recombinant Helicobacter pylori urease B subunit as referenced above; "CT" means cholera toxin; and "HAP" means hydroxyapatite crystals. "sacrifice" means the date of sacrifice as measured from the last immunization at day 44. "N.D." indicates that no data were available. The results of the urease test performed from biopsies of the stomach of each animal are expressed as O.D. values at 550 nm. The infection is rated as positive (indicated by a "+") where the O.D. value at 550 nm is over twice the standard deviation found in the mature mice (0.04 O.D.). The

1095539-091891
T08160-6EY55660

background value subtracted from the O.D. value for each mouse was 0.04 O.D. In Table 10, "% cleared" indicates clearance of infection as measured by the urease assay.

It will be seen from the results set out in Tables 9 and 10, and in Figure 8, that administration of the Helicobacter pylori ure B subunit to mice infected with H. felis resulted in the clearance of infection in eight out of ten mice as assayed when the mice were sacrificed 2.5 weeks after the last immunization. When the assay was performed when the mice were sacrificed eight weeks after the last immunization, the infection was cleared in six out of nine mice. If the higher value of 0.2 is chosen to define infection as referred to in the discussion of Table 8, under this method of interpretation, treatment with the ure B subunit resulted in clearance of infection in nine out of ten mice sacrificed at 2.5 weeks.

Table 10 indicates that statistically significant therapeutic treatment of H. felis infection is obtained with oral immunization using recombinant Helicobacter pylori urease B subunit as compared to that obtained by sham immunization with cholera toxin and hydroxyapatite alone. Table 10 shows that, when measured at 2.5 weeks after the last immunization, 80% of the infected mice were cleared of infection. When measured at eight weeks after the last immunization, 67% of the infected mice were cleared of infection. None of the sham immunized mice were

0955739-09101
105T60-6575560

cleared of infection when measured at either 2.5 or eight weeks after the last immunization. The significance of the reduction in the percentage of mice cleared of infection when measured at eight weeks as compared to 2.5 weeks after the last immunization is unclear at present.

TABLE 11

<u>Mouse No.</u>	<u>Urease Test</u>	<u>Total anti-ureB Igs in Blood</u>	<u>Total IgAs in Feces</u>	<u>ureB IgA's in Feces</u>	<u>ure B IgA's in Feces/ Total IgA's in Feces</u>
<u>Ex. 4:</u> 8	0.51	0.27	0.77	0.05	0.06
9	0.48	0.23	0.65	0.02	0.03
10	0.61	0.25	0.43	0.08	0.18
11	0.44	0.36	0.69	0.03	0.04
12	0.51	0.31	0.58	0.00	0.00
13	0.48	0.32	0.67	0.11	0.17
14	0.44	0.27	0.51	0.03	0.05
15	0.58	0.29	0.82	0.07	0.08
16	0.48	0.30	0.77	0.18	0.23
17	0.53	0.30	0.53	0.09	0.18
18	0.36	0.22	0.51	0.00	0.00
19	0.01	0.47	0.62	0.47	0.75
20	0.43	0.35	0.57	0.15	0.26
21	0.07	0.25	0.64	0.11	0.17

<u>Mouse No.</u>	<u>Urease Test</u>	<u>Total anti-ureB Igs in Blood</u>	<u>Total IgAs in Feces</u>	<u>ureB IgA's in Feces</u>	<u>ure B IgA's in Feces/ Total IgA's in Feces</u>
22	0.39	0.27	0.50	0.15	0.30
23	0.10	0.27	1.08	ND	ND
24	0.38	0.28	0.66	ND	ND
25	0.06	0.28	0.53	ND	ND
Ex.5: 7	0.02	0.11	0.61	0.06	0.11
8	0.06	0.13	0.62	0.02	0.03
9	0.01	0.15	0.62	0.04	0.07
10	0.00	0.15	0.59	0.03	0.05
11	0.02	0.19	0.60	0.15	0.24
12	0.02	0.16	0.60	0.00	-0.01
13	0.02	0.19	0.66	0.07	0.11
14	0.44	0.18	0.62	0.00	0.00
15	0.00	0.10	0.64	0.08	0.12
16	0.03	0.14	0.62	0.29	0.46
25	0.26	0.35	0.98	0.07	0.07
26	0.30	0.38	1.12	0.04	0.03
27	0.00	0.39	1.15	0.01	0.01
28	0.17	0.33	0.97	0.02	0.02
29	N.D.*	0.38	1.00	0.08	0.08
30	0.00	0.34	0.69	0.04	0.06
31	0.00	0.38	0.76	0.08	0.10
32	0.00	0.37	0.58	0.02	0.04
33	0.00	0.38	1.07	0.22	0.21

<u>Mouse No.</u>	<u>Urease Test</u>	<u>Total anti-ureB Igs in Blood</u>	<u>Total IgAs in Feces</u>	<u>ureB IgA's in Feces</u>	<u>ure B IgA's in Feces/ Total IgA's in Feces</u>
34	0.00	0.35	0.64	0.06	0.09

In Table 11, the mice that were treated with ureB, used in Examples 4 and 5, were evaluated for the antibody content of the blood and the feces. The mouse numbers used correspond to the numbers used in Tables 8 and 9. The urease test results correspond to the results reported in Tables 8 and 9. "ND*" indicates that no data was available because one of the stomachs, from mice numbered 18-21, was lost. The inventors are unable to determine which of the sets of data from mice numbered 18-21 is incomplete. Therefore, the "ND" is not meant to be specifically assigned to mouse number 21.

In the assays of the mice from Example 4, and the mice numbered 1-10 from Example 5, the inventors had difficulty obtaining adequate samples for analysis.

The clearance of H. felis infection by oral administration of the Helicobacter pylori ure B subunit was not expected and is therefore novel. The results described herein also teach that the ure B subunit can be used to treat Helicobacter infection.

One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The urease peptides, mucosal adjuvants, carriers, and antibodies, along with the methods, procedures, treatments, and assays, are exemplary and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention as defined by the scope of the claims.

108T60-6E755660